

TABLE 12. Estimated percent of children allowed to carry medication in school among children¹ with current² asthma status who attended school in the past year³ by state/territory - BRFSS Asthma Call-back Survey, United States, 2006-2010

State/Territory	Current			
	Sample size [±]	%	SE [§]	95% CI [†]
Arizona	96	27.4	6.5	(14.7 - 40.1)
California	145	52.4	5.5	(41.6 - 63.2)
Connecticut	305	37.7	3.7	(30.4 - 45.0)
District of Columbia	95	46.3	6.4	(33.8 - 58.8)
Georgia	289	62.5	3.9	(54.9 - 70.1)
Hawaii	313	59.8	3.6	(52.7 - 66.9)
Illinois	154	66.4	4.8	(57.0 - 75.8)
Indiana	289	50.3	3.8	(42.9 - 57.7)
Iowa	208	65.7	3.9	(58.1 - 73.3)
Kansas	497	44.9	2.7	(39.6 - 50.2)
Louisiana	86	40.5	6.3	(28.2 - 52.8)
Maine	242	55.2	3.8	(47.8 - 62.6)
Maryland	311	39.7	3.6	(32.6 - 46.8)
Massachusetts	229	39.6	4.5	(30.8 - 48.4)
Michigan	403	54.4	3.2	(48.1 - 60.7)
Mississippi	58	54.6	8.7	(37.5 - 71.7)
Missouri	199	40.7	5.0	(30.9 - 50.5)
Montana	184	65.4	4.7	(56.2 - 74.6)
Nebraska	404	47.3	4.0	(39.5 - 55.1)
New Hampshire	126	58.0	5.0	(48.2 - 67.8)
New Jersey	263	45.7	4.1	(37.7 - 53.7)
New Mexico	206	61.5	4.4	(52.9 - 70.1)
New York	238	45.3	4.3	(36.9 - 53.7)
North Dakota	86	77.3	5.3	(66.9 - 87.7)
Ohio	184	57.3	5.6	(46.3 - 68.3)
Oklahoma	206	55.4	4.4	(46.8 - 64.0)
Oregon	161	44.9	4.8	(35.5 - 54.3)
Pennsylvania	109	47.2	6.6	(34.3 - 60.1)
Rhode Island	139	47.7	5.5	(36.9 - 58.5)
Texas	409	45.9	4.0	(38.1 - 53.7)
Utah	291	72.9	3.5	(66.0 - 79.8)
Vermont	320	48.6	3.3	(42.1 - 55.1)
Virginia	70	28.2	5.9	(16.6 - 39.8)
Washington	284	60.3	3.7	(53.0 - 67.6)
West Virginia	158	40.1	4.4	(31.5 - 48.7)
Wisconsin	122	62.5	7.0	(48.8 - 76.2)
36 state total	7,879	49.8	1.2	(47.4 - 52.2)
Median		49.5		
Range		27.4-77.3		

¹ Aged < 18 years

² "Yes" response to "Do you still have asthma?"

³ Attended school outside the home. Home Schooled are excluded.

[±] Sample size is for multiple years of participation. Responses of "Don't Know" and "Refused" are excluded.

[§] Standard error

[†] Confidence interval