

TABLE 3a. Estimated percent for time since first diagnosis among children¹ with current² asthma by state/territory - BRFSS Asthma Call-back Survey, United States, 2006-2010

State/Territory	Sample size [±]	5 or fewer years				More than 5 years			
		Sample size	%	SE [§]	95% CI [†]	Sample size	%	SE	95% CI
Arizona	126	54	52.1	7.1	(38.2 - 66.0)	72	47.9	7.1	(34.0 - 61.8)
California	178	88	50.8	5.0	(41.0 - 60.6)	90	49.2	5.0	(39.4 - 59.0)
Connecticut	357	176	49.3	3.5	(42.4 - 56.2)	181	50.7	3.5	(43.8 - 57.6)
District of Columbia	122	54	40.4	5.6	(29.4 - 51.4)	68	59.6	5.6	(48.6 - 70.6)
Georgia	358	156	49.7	3.7	(42.4 - 57.0)	202	50.3	3.7	(43.0 - 57.6)
Hawaii	427	178	46.6	3.1	(40.5 - 52.7)	249	53.4	3.1	(47.3 - 59.5)
Illinois	172	75	48.6	4.9	(39.0 - 58.2)	97	51.4	4.9	(41.8 - 61.0)
Indiana	354	177	55.8	3.4	(49.1 - 62.5)	177	44.2	3.4	(37.5 - 50.9)
Iowa	253	136	57.6	3.6	(50.5 - 64.7)	117	42.4	3.6	(35.3 - 49.5)
Kansas	603	289	53.1	2.4	(48.4 - 57.8)	314	46.9	2.4	(42.2 - 51.6)
Louisiana	107	42	45.6	5.9	(34.0 - 57.2)	65	54.4	5.9	(42.8 - 66.0)
Maine	294	131	46.4	3.3	(39.9 - 52.9)	163	53.6	3.3	(47.1 - 60.1)
Maryland	394	177	48.3	3.3	(41.8 - 54.8)	217	51.7	3.3	(45.2 - 58.2)
Massachusetts	276	117	48.5	4.6	(39.5 - 57.5)	159	51.5	4.6	(42.5 - 60.5)
Michigan	492	232	50.7	3.0	(44.8 - 56.6)	260	49.3	3.0	(43.4 - 55.2)
Mississippi	81	38	59.6	6.6	(46.7 - 72.5)	43	40.4	6.6	(27.5 - 53.3)
Missouri	234	116	50.4	4.8	(41.0 - 59.8)	118	49.6	4.8	(40.2 - 59.0)
Montana	214	106	53.7	4.1	(45.7 - 61.7)	108	46.3	4.1	(38.3 - 54.3)
Nebraska	487	242	51.7	3.9	(44.1 - 59.3)	245	48.3	3.9	(40.7 - 55.9)
New Hampshire	156	76	50.8	4.8	(41.4 - 60.2)	80	49.2	4.8	(39.8 - 58.6)
New Jersey	306	139	43.1	3.8	(35.7 - 50.5)	167	56.9	3.8	(49.5 - 64.3)
New Mexico	240	95	47.7	4.3	(39.3 - 56.1)	145	52.3	4.3	(43.9 - 60.7)
New York	287	130	50.7	3.9	(43.1 - 58.3)	157	49.3	3.9	(41.7 - 56.9)
North Dakota	102	51	57.4	5.7	(46.2 - 68.6)	51	42.6	5.7	(31.4 - 53.8)
Ohio	229	121	56.3	5.1	(46.3 - 66.3)	108	43.7	5.1	(33.7 - 53.7)
Oklahoma	254	125	53.6	3.9	(46.0 - 61.2)	129	46.4	3.9	(38.8 - 54.0)
Oregon	198	91	52.1	4.3	(43.7 - 60.5)	107	47.9	4.3	(39.5 - 56.3)
Pennsylvania	135	64	49.0	5.7	(37.8 - 60.2)	71	51.0	5.7	(39.8 - 62.2)
Rhode Island	161	73	52.5	4.5	(43.7 - 61.3)	88	47.5	4.5	(38.7 - 56.3)
Texas	495	226	55.3	3.7	(48.0 - 62.6)	269	44.7	3.7	(37.4 - 52.0)
Utah	380	194	53.4	3.4	(46.7 - 60.1)	186	46.6	3.4	(39.9 - 53.3)
Vermont	395	173	45.4	3.0	(39.5 - 51.3)	222	54.6	3.0	(48.7 - 60.5)
Virginia	88	42	49.7	8.6	(32.8 - 66.6)	46	50.3	8.6	(33.4 - 67.2)
Washington	349	164	52.2	3.4	(45.5 - 58.9)	185	47.8	3.4	(41.1 - 54.5)
West Virginia	204	88	48.0	4.1	(40.0 - 56.0)	116	52.0	4.1	(44.0 - 60.0)
Wisconsin	148	68	58.6	5.4	(48.0 - 69.2)	80	41.4	5.4	(30.8 - 52.0)
36 state total	9,656	4,504	51.2	1.1	(49.0 - 53.4)	5,152	48.8	1.1	(46.6 - 51.0)
Median			50.8				49.3		
Range			40.4-59.6				40.4-59.6		

¹ Age < 18 years

² "Yes" response to "Do you still have asthma?"

[±] Sample size is for multiple years of participation. Responses of "Don't Know" and "Refused" are excluded.

[§] Standard error

[†] Confidence interval