

TABLE 4b. Estimated percent reporting an asthma attack or episode in the past year among children¹ with active² asthma by state/territory - BRFSS Asthma Call-back Survey, United States, 2006-2010

State/Territory	Sample size [±]	Attack in past year				No attack in past year				No symptoms in past year			
		Sample size	%	SE [§]	95% CI [†]	Sample size	%	SE	95% CI	Sample size	%	SE	95% CI
Arizona	150	84	57.1	6.5	(44.4 - 69.8)	51	32.6	6.3	(20.3 - 44.9)	15	N/A	N/A	(N/A - N/A)
California	214	116	55.8	4.6	(46.8 - 64.8)	66	31.4	4.4	(22.8 - 40.0)	32	12.8	2.9	(7.1 - 18.5)
Connecticut	427	222	51.2	3.1	(45.1 - 57.3)	130	28.1	2.8	(22.6 - 33.6)	75	20.7	2.7	(15.4 - 26.0)
District of Columbia	148	72	49.3	5.1	(39.3 - 59.3)	49	31.2	4.6	(22.2 - 40.2)	27	19.5	4.1	(11.5 - 27.5)
Georgia	414	223	50.2	3.5	(43.3 - 57.1)	129	34.9	3.4	(28.2 - 41.6)	62	14.9	2.4	(10.2 - 19.6)
Hawaii	453	205	45.8	3.0	(39.9 - 51.7)	168	37.1	2.9	(31.4 - 42.8)	80	17.1	2.2	(12.8 - 21.4)
Illinois	200	101	53.4	4.5	(44.6 - 62.2)	63	29.2	4.1	(21.2 - 37.2)	36	17.4	3.2	(11.1 - 23.7)
Indiana	394	222	55.7	3.4	(49.0 - 62.4)	128	33.8	3.2	(27.5 - 40.1)	44	10.5	2.0	(6.6 - 14.4)
Iowa	280	134	50.8	3.5	(43.9 - 57.7)	96	35.1	3.3	(28.6 - 41.6)	50	14.1	2.3	(9.6 - 18.6)
Kansas	667	335	50.6	2.3	(46.1 - 55.1)	231	33.2	2.1	(29.1 - 37.3)	101	16.1	1.8	(12.6 - 19.6)
Louisiana	121	60	48.7	5.5	(37.9 - 59.5)	45	42.7	5.5	(31.9 - 53.5)	16	8.6	2.3	(4.1 - 13.1)
Maine	344	172	51.7	3.1	(45.6 - 57.8)	118	31.9	2.9	(26.2 - 37.6)	54	16.5	2.4	(11.8 - 21.2)
Maryland	477	254	52.6	3.0	(46.7 - 58.5)	134	29.6	2.7	(24.3 - 34.9)	89	17.8	2.3	(13.3 - 22.3)
Massachusetts	320	150	48.5	4.0	(40.7 - 56.3)	118	34.6	3.7	(27.3 - 41.9)	52	16.9	3.4	(10.2 - 23.6)
Michigan	542	299	56.3	2.8	(50.8 - 61.8)	145	27.2	2.5	(22.3 - 32.1)	98	16.5	2.0	(12.6 - 20.4)
Mississippi	95	53	54.8	6.7	(41.7 - 67.9)	34	38.4	6.7	(25.3 - 51.5)	8	N/A	N/A	(N/A - N/A)
Missouri	259	140	52.8	4.5	(44.0 - 61.6)	89	34.2	4.2	(26.0 - 42.4)	30	13.0	3.7	(5.7 - 20.3)
Montana	258	139	56.4	3.8	(49.0 - 63.8)	78	29.1	3.5	(22.2 - 36.0)	41	14.5	2.6	(9.4 - 19.6)
Nebraska	547	260	50.5	3.6	(43.4 - 57.6)	191	33.2	3.5	(26.3 - 40.1)	96	16.3	2.8	(10.8 - 21.8)
New Hampshire	177	93	53.0	4.5	(44.2 - 61.8)	57	32.7	4.4	(24.1 - 41.3)	27	14.3	3.4	(7.6 - 21.0)
New Jersey	346	157	43.1	3.8	(35.7 - 50.5)	124	37.0	3.8	(29.6 - 44.4)	65	19.9	3.2	(13.6 - 26.2)
New Mexico	268	130	52.8	4.2	(44.6 - 61.0)	88	29.6	3.6	(22.5 - 36.7)	50	17.6	3.2	(11.3 - 23.9)
New York	346	172	51.8	3.8	(44.4 - 59.2)	115	30.4	3.5	(23.5 - 37.3)	59	17.8	3.2	(11.5 - 24.1)
North Dakota	124	62	50.0	5.4	(39.4 - 60.6)	44	36.8	5.2	(26.6 - 47.0)	18	13.2	3.3	(6.7 - 19.7)
Ohio	266	126	48.7	4.5	(39.9 - 57.5)	94	35.7	4.5	(26.9 - 44.5)	46	15.5	2.9	(9.8 - 21.2)
Oklahoma	261	144	57.9	3.8	(50.5 - 65.3)	84	31.4	3.6	(24.3 - 38.5)	33	10.7	2.1	(6.6 - 14.8)
Oregon	226	109	55.3	4.1	(47.3 - 63.3)	67	27.6	3.6	(20.5 - 34.7)	50	17.0	2.6	(11.9 - 22.1)
Pennsylvania	155	77	53.0	5.5	(42.2 - 63.8)	50	32.0	4.9	(22.4 - 41.6)	28	15.0	3.7	(7.7 - 22.3)
Rhode Island	189	101	54.2	4.2	(46.0 - 62.4)	48	25.2	3.8	(17.8 - 32.6)	40	20.6	3.3	(14.1 - 27.1)
Texas	572	306	52.1	3.5	(45.2 - 59.0)	180	33.4	3.5	(26.5 - 40.3)	86	14.5	2.6	(9.4 - 19.6)
Utah	422	257	63.1	3.4	(56.4 - 69.8)	113	25.6	3.1	(19.5 - 31.7)	52	11.3	1.9	(7.6 - 15.0)
Vermont	449	216	51.1	2.9	(45.4 - 56.8)	162	33.3	2.6	(28.2 - 38.4)	71	15.6	2.0	(11.7 - 19.5)
Virginia	101	53	58.2	8.2	(42.1 - 74.3)	31	30.4	8.2	(14.3 - 46.5)	17	11.4	2.9	(5.7 - 17.1)
Washington	386	196	49.3	3.3	(42.8 - 55.8)	125	33.8	3.1	(27.7 - 39.9)	65	16.9	2.7	(11.6 - 22.2)
West Virginia	224	116	53.8	3.9	(46.2 - 61.4)	84	37.6	3.7	(30.3 - 44.9)	24	8.6	1.9	(4.9 - 12.3)
Wisconsin	200	102	54.3	4.6	(45.3 - 63.3)	69	32.8	4.3	(24.4 - 41.2)	29	12.9	2.9	(7.2 - 18.6)
36 state total	11,022	5,658	52.9	1.1	(50.7 - 55.1)	3,598	32.3	1.0	(30.3 - 34.3)	1,766	14.8	0.7	(13.4 - 16.2)
Median			52.7				32.8				15.6		
Range			43.1-63.1				25.2-42.7				8.6-20.7		

¹ Aged < 18 years

² Reported any of the following during the past year: asthma medication use, asthma symptoms, or MD visits for asthma

[±] Sample size is for multiple years of participation. Responses of "Don't Know" and "Refused" are excluded.

[§] Standard error

[†] Confidence interval

N/A: Estimate not available (N/A) if the unweighted sample size for the denominator was < 50 or if the relative standard error was > 0.30