

TABLE 6. Estimated percent receiving an asthma management plan among children¹ with current² or active³ asthma status by state/territory - BRFSS Asthma Call-back Survey, United States, 2006-2010

State/Territory	Current				Active			
	Sample size [±]	%	SE [§]	95% CI [†]	Sample size	%	SE	95% CI
Arizona	124	45.4	7.2	(31.3 - 59.5)	148	44.9	6.4	(32.4 - 57.4)
California	175	46.2	4.9	(36.6 - 55.8)	212	43.8	4.6	(34.8 - 52.8)
Connecticut	351	52.5	3.6	(45.4 - 59.6)	420	51.1	3.3	(44.6 - 57.6)
District of Columbia	120	53.7	5.8	(42.3 - 65.1)	146	51.5	5.3	(41.1 - 61.9)
Georgia	350	45.6	3.6	(38.5 - 52.7)	410	43.7	3.3	(37.2 - 50.2)
Hawaii	421	36.7	3.1	(30.6 - 42.8)	447	35.2	2.9	(29.5 - 40.9)
Illinois	172	49.8	5.0	(40.0 - 59.6)	200	50.5	4.6	(41.5 - 59.5)
Indiana	343	47.7	3.6	(40.6 - 54.8)	386	44.7	3.3	(38.2 - 51.2)
Iowa	252	30.9	3.4	(24.2 - 37.6)	278	31.7	3.1	(25.6 - 37.8)
Kansas	593	47.2	2.5	(42.3 - 52.1)	661	46.9	2.3	(42.4 - 51.4)
Louisiana	106	51.3	6.0	(39.5 - 63.1)	120	50.3	5.6	(39.3 - 61.3)
Maine	283	61.5	3.3	(55.0 - 68.0)	332	59.8	3.1	(53.7 - 65.9)
Maryland	387	45.7	3.4	(39.0 - 52.4)	469	46.3	3.1	(40.2 - 52.4)
Massachusetts	269	49.6	4.4	(41.0 - 58.2)	314	49.0	4.1	(41.0 - 57.0)
Michigan	484	43.7	3.0	(37.8 - 49.6)	540	43.2	2.8	(37.7 - 48.7)
Mississippi	80	28.4	5.9	(16.8 - 40.0)	94	27.4	5.5	(16.6 - 38.2)
Missouri	231	50.8	4.8	(41.4 - 60.2)	255	49.9	4.6	(40.9 - 58.9)
Montana	212	44.9	4.3	(36.5 - 53.3)	253	46.9	4.0	(39.1 - 54.7)
Nebraska	481	46.7	3.9	(39.1 - 54.3)	544	44.2	3.6	(37.1 - 51.3)
New Hampshire	145	47.6	5.0	(37.8 - 57.4)	169	45.4	4.6	(36.4 - 54.4)
New Jersey	304	52.8	4.0	(45.0 - 60.6)	344	49.2	3.9	(41.6 - 56.8)
New Mexico	238	42.4	4.2	(34.2 - 50.6)	266	41.9	4.0	(34.1 - 49.7)
New York	280	43.8	4.2	(35.6 - 52.0)	339	38.9	3.8	(31.5 - 46.3)
North Dakota	97	43.4	6.1	(31.4 - 55.4)	119	42.4	5.5	(31.6 - 53.2)
Ohio	224	48.7	4.8	(39.3 - 58.1)	259	52.4	4.8	(43.0 - 61.8)
Oklahoma	252	45.5	3.9	(37.9 - 53.1)	257	46.2	3.9	(38.6 - 53.8)
Oregon	192	31.6	4.3	(23.2 - 40.0)	222	32.4	4.0	(24.6 - 40.2)
Pennsylvania	133	44.1	5.8	(32.7 - 55.5)	156	43.7	5.4	(33.1 - 54.3)
Rhode Island	158	43.6	4.7	(34.4 - 52.8)	184	45.4	4.3	(37.0 - 53.8)
Texas	482	47.1	3.7	(39.8 - 54.4)	562	45.7	3.5	(38.8 - 52.6)
Utah	375	43.2	3.4	(36.5 - 49.9)	420	41.7	3.2	(35.4 - 48.0)
Vermont	384	51.8	3.0	(45.9 - 57.7)	437	50.2	2.9	(44.5 - 55.9)
Virginia	84	45.7	8.8	(28.5 - 62.9)	96	46.3	8.1	(30.4 - 62.2)
Washington	337	36.4	3.4	(29.7 - 43.1)	382	36.0	3.3	(29.5 - 42.5)
West Virginia	194	30.2	3.8	(22.8 - 37.6)	215	27.9	3.6	(20.8 - 35.0)
Wisconsin	147	46.9	5.8	(35.5 - 58.3)	206	45.4	4.8	(36.0 - 54.8)
36 state total	9,460	45.9	1.1	(43.7 - 48.1)	10,862	44.6	1.1	(42.4 - 46.8)
Median		45.7				45.4		
Range		28.4-61.5				27.4-59.8		

¹ Aged < 18 years

² "Yes" response to "Do you still have asthma?"

³ Reported any of the following during the past year: asthma medication use, asthma symptoms, or MD visits for asthma

[±] Sample size is for multiple years of participation. Responses of "Don't Know" and "Refused" are excluded.

[§] Standard error

[†] Confidence interval