

**Table L1**  
**Child Lifetime Asthma Prevalence Rate (Percent)**  
**and Prevalence (Number) by State or Territory: BRFSS 2007**

State	Sample Size	Prevalence (Percent)	Standard Error	95% CI * (Percent)	    	Prevalence (Number)	95% CI * (Number)
Total **	59,849	13.5	0.27	(13.0 - 14.1)		7,532,693	(7,228,449 - 7,836,936)
AK	942	11.0	1.48	(8.4 - 14.2)		20,830	(15,261 - 26,400)
AZ	1,258	13.9	1.71	(10.9 - 17.6)		224,825	(169,120 - 280,530)
CA	1,687	13.2	0.97	(11.4 - 15.2)		1,289,140	(1,103,150 - 1,475,130)
CT	1,940	14.7	0.98	(12.9 - 16.7)		120,988	(104,933 - 137,042)
DC	798	19.7	1.98	(16.1 - 23.9)		22,522	(17,757 - 27,286)
GA	2,092	15.3	1.04	(13.4 - 17.5)		362,710	(313,423 - 411,996)
HI	2,145	18.2	1.02	(16.3 - 20.3)		55,478	(49,296 - 61,660)
IL	1,488	11.3	1.05	(9.4 - 13.5)		363,468	(295,124 - 431,812)
IN	1,599	11.3	0.96	(9.5 - 13.3)		179,341	(149,412 - 209,269)
IA	1,509	8.7	0.78	(7.3 - 10.4)		57,762	(47,551 - 67,973)
KS	2,381	11.2	0.73	(9.9 - 12.7)		75,715	(65,943 - 85,488)
ME	1,588	13.3	0.97	(11.5 - 15.3)		36,508	(31,261 - 41,755)
MD	2,384	13.6	0.95	(11.9 - 15.6)		190,043	(163,564 - 216,521)
MA	1,186	14.6	1.31	(12.2 - 17.4)		209,137	(172,057 - 246,216)
MI	2,005	14.2	1.00	(12.3 - 16.3)		350,245	(300,140 - 400,350)
MS	2,169	14.8	0.97	(13.0 - 16.8)		109,587	(95,207 - 123,966)
MO	1,362	13.1	1.11	(11.0 - 15.4)		180,576	(151,240 - 209,911)
MT	1,516	9.8	0.92	(8.1 - 11.7)		20,087	(16,348 - 23,827)
NE	3,075	11.1	1.01	(9.3 - 13.3)		48,234	(39,640 - 56,828)
NV	1,169	10.3	1.09	(8.3 - 12.6)		65,582	(51,995 - 79,168)
NH	1,517	11.4	0.94	(9.7 - 13.4)		34,002	(28,412 - 39,592)
NJ	913	14.5	1.53	(11.8 - 17.8)		310,559	(246,372 - 374,745)
NM	1,828	13.0	0.98	(11.2 - 15.1)		63,854	(54,327 - 73,380)
NY	1,563	16.1	1.24	(13.9 - 18.7)		718,245	(604,852 - 831,639)
OH	2,865	13.7	0.78	(12.2 - 15.3)		371,762	(329,462 - 414,062)
OK	2,046	15.5	0.94	(13.8 - 17.4)		132,346	(116,498 - 148,194)

Notes:  
\* CI denotes confidence interval.  
\*\* Total includes data from 33 states plus the District of Columbia and excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

**Table L1**  
**Child Lifetime Asthma Prevalence Rate (Percent)**  
**and Prevalence (Number) by State or Territory: BRFSS 2007**

State	Sample Size	Prevalence (Percent)	Standard Error	95% CI * (Percent)	    	Prevalence (Number)	95% CI * (Number)
OR	1,241	11.9	1.06	(10.0 - 14.1)		100,186	(82,393 - 117,978)
PA	3,130	14.7	1.03	(12.8 - 16.9)		406,319	(349,404 - 463,235)
TX	2,590	13.7	0.95	(12.0 - 15.7)		877,720	(755,937 - 999,502)
UT	2,003	11.7	0.90	(10.0 - 13.5)		88,037	(74,357 - 101,716)
VT	1,756	12.4	0.89	(10.7 - 14.2)		16,118	(13,802 - 18,434)
VA	1,649	13.4	1.15	(11.2 - 15.8)		244,053	(201,971 - 286,135)
WV	1,062	13.7	1.19	(11.5 - 16.2)		51,331	(42,493 - 60,169)
WI	1,393	10.6	1.12	(8.6 - 13.0)		135,388	(107,091 - 163,684)
Territories							
PR	1,070	28.5	1.59	(25.5 - 31.7)		296,304	(262,424 - 330,184)

Notes:  
\* CI denotes confidence interval.  
\*\* Total includes data from 33 states plus the District of Columbia and excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.