

Table C2_1
Child Current Asthma Prevalence Rate (Percent)
by Sex and State or Territory: BRFSS 2008

State	Male Sample Size	Male Prevalence	Male Standard Error	Male 95% CI *	Female Sample Size	Female Prevalence	Female Standard Error	Female 95% CI *
Total **	39,407	10.4	0.30	(9.9 - 11.0)	36,499	7.6	0.25	(7.1 - 8.1)
AZ	767	10.5	2.26	(6.8 - 15.9)	736	5.7	1.59	(3.2 - 9.7)
CA	1,899	8.8	0.81	(7.4 - 10.5)	1,759	5.5	0.67	(4.3 - 7.0)
CT	854	11.3	1.49	(8.7 - 14.6)	765	10.9	1.66	(8.0 - 14.6)
DC	468	14.9	2.34	(10.9 - 20.1)	419	10.4	2.17	(6.9 - 15.5)
GA	868	13.6	1.79	(10.5 - 17.5)	782	10.4	1.46	(7.8 - 13.6)
HI	1,053	15.6	1.39	(13.0 - 18.5)	957	9.7	1.26	(7.5 - 12.5)
ID	809	8.2	1.29	(6.0 - 11.2)	787	5.0	1.01	(3.3 - 7.4)
IL	753	9.4	1.33	(7.1 - 12.4)	786	7.5	1.19	(5.5 - 10.2)
IN	716	9.6	1.64	(6.8 - 13.3)	679	7.4	1.14	(5.5 - 10.0)
IA	890	6.6	0.91	(5.0 - 8.6)	796	6.7	0.98	(5.0 - 8.9)
KS	1,215	10.1	0.97	(8.3 - 12.1)	1,196	7.9	0.87	(6.3 - 9.8)
KY	1,090	10.0	1.23	(7.9 - 12.7)	1,077	8.9	1.13	(6.9 - 11.4)
ME	827	10.4	1.14	(8.3 - 12.8)	815	8.4	1.04	(6.6 - 10.7)
MD	1,462	10.9	1.05	(9.0 - 13.2)	1,370	8.2	1.02	(6.4 - 10.5)
MA	971	12.4	1.54	(9.6 - 15.7)	837	7.4	1.37	(5.1 - 10.6)
MI	1,283	12.5	1.40	(10.0 - 15.5)	1,163	9.0	0.94	(7.3 - 11.0)
MS	963	13.2	1.44	(10.6 - 16.3)	919	7.3	1.16	(5.3 - 9.9)
MO	674	12.0	1.81	(8.9 - 16.0)	664	8.2	1.43	(5.8 - 11.5)
MT	881	7.4	1.15	(5.4 - 10.0)	785	6.1	1.10	(4.3 - 8.7)
NE	2,272	7.3	0.97	(5.6 - 9.4)	2,214	5.0	0.76	(3.7 - 6.7)
NV	741	7.8	1.34	(5.5 - 10.9)	621	7.0	1.67	(4.4 - 11.1)
NH	938	10.5	1.26	(8.3 - 13.3)	865	6.7	0.97	(5.0 - 8.8)
NJ	1,762	10.3	0.96	(8.5 - 12.3)	1,570	6.8	0.83	(5.3 - 8.6)
NM	985	9.3	1.20	(7.2 - 11.9)	865	5.9	0.88	(4.4 - 7.9)
NY	1,025	10.8	1.19	(8.7 - 13.4)	988	10.5	1.38	(8.1 - 13.6)
ND	647	7.2	1.25	(5.1 - 10.0)	639	3.7	0.72	(2.5 - 5.4)

Notes:
* CI denotes confidence interval.
** Total includes data from 37 states plus the District of Columbia and excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

Table C2_1
Child Current Asthma Prevalence Rate (Percent)
by Sex and State or Territory: BRFSS 2008

State	Male Sample Size	Male Prevalence	Male Standard Error	Male 95% CI *		Female Sample Size	Female Prevalence	Female Standard Error	Female 95% CI *
OH	1,677	9.9	1.01	(8.1 - 12.1)		1,555	8.9	0.95	(7.2 - 10.9)
OK	1,196	11.9	1.15	(9.8 - 14.3)		1,029	8.1	0.97	(6.4 - 10.3)
OR	613	5.3	1.05	(3.6 - 7.8)		542	6.0	1.24	(3.9 - 8.9)
PA	1,619	11.8	1.37	(9.3 - 14.7)		1,528	7.8	1.01	(6.1 - 10.0)
RI	677	15.3	1.61	(12.4 - 18.7)		588	9.6	1.50	(7.0 - 12.9)
TX	1,633	11.6	1.25	(9.3 - 14.2)		1,554	6.9	0.93	(5.3 - 9.0)
UT	1,254	7.1	0.93	(5.5 - 9.2)		987	6.2	1.09	(4.4 - 8.7)
VT	907	12.8	1.32	(10.4 - 15.6)		836	7.0	0.98	(5.3 - 9.2)
VA	675	10.0	1.68	(7.2 - 13.8)		652	9.3	1.54	(6.7 - 12.8)
WV	470	13.2	1.85	(10.0 - 17.3)		484	9.7	1.52	(7.1 - 13.1)
WI	711	9.1	1.60	(6.4 - 12.7)		657	5.9	1.21	(3.9 - 8.7)
WY	1,162	7.5	0.85	(6.0 - 9.4)		1,033	4.4	0.72	(3.2 - 6.1)
Territories									
PR	578	16.8	1.95	(13.3 - 21.0)		537	11.0	1.53	(8.3 - 14.3)

Notes:
* CI denotes confidence interval.
** Total includes data from 37 states plus the District of Columbia and excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.