Table L4
Child Lifetime Asthma Prevalence
and Weighted Number by Race and State or Territory: BRFSS 2018

State	Race	Unweighted Number	Prevalence (Percent)	Standard Error	95% CI * (Percent)		Weighted Number	95% CI* Weighted Number
Total **	White	40,115	10.2	0.29	(9.7–10.8)	||	2,979,575	(2,807,250–3,151,899)
Total **	Black	5,085	16.3	0.88	(14.6–18.1)	||	1,158,035	(1,027,779–1,288,291)
Total **	Other	8,838	11.2	0.63	(10.1–12.5)	||	1,039,580	(921,115–1,158,045)
CA	White	1,899	9.9	0.82	(8.5–11.7)	||	541,605	(451,934–631,276)
CA	Black	172	16.7	3.63	(10.7–25.1)	||	103,846	(55,473–152,219)
CA	Other	798	10.7	1.42	(8.2–13.8)	||	294,920	(214,237–375,603)
CT	White	1,465	14.0	1.23	(11.7–16.6)	||	57,187	(46,861–67,514)
CT	Black	216	16.0	2.72	(11.4–22.1)	||	16,849	(11,033–22,665)
CT	Other	308	17.0	2.64	(12.5–22.9)	||	22,556	(15,134–29,977)
DC	White	286	6.5	1.56	(4.0–10.3)	||	2,465	(1,285–3,644)
DC	Black	349	23.0	2.71	(18.1–28.8)	||	14,025	(10,521–17,529)
DC	Other	79	15.2	4.58	(8.2–26.5)	||	2,697	(983–4,410)
FL	White	2,063	11.5	1.33	(9.2–14.4)	||	286,610	(219,436–353,785)
FL	Black	406	15.5	3.23	(10.2–23.0)	||	148,235	(82,575–213,896)
FL	Other	388	12.0	3.07	(7.1–19.4)	||	74,842	(35,089–114,595)
GA	White	1,157	11.1	1.10	(9.1–13.4)	||	128,733	(103,000–154,465)
GA	Black	564	14.3	1.71	(11.3–18.0)	||	122,390	(92,379–152,402)
GA	Other	339	11.0	2.06	(7.6–15.7)	||	37,130	(23,066–51,194)
HI	White	329	13.2	2.74	(8.7–19.6)	||	6,684	(3,730–9,639)
HI	Black	14	35.9	18.02	(10.8–72.2)	||	1,424	(0–3,202)
HI	Other	1,493	18.3	1.25	(16.0–20.9)	||	44,045	(37,891–50,199)
IL	White	872	8.8	1.10	(6.9–11.2)	||	161,315	(121,607–201,023)
IL	Black	181	11.0	2.64	(6.7–17.3)	||	53,716	(27,430–80,002)
IL	Other	199	9.7	2.65	(5.6–16.2)	||	39,461	(17,195–61,727)
IN	White	1,125	10.9	1.11	(8.9–13.3)	||	105,236	(83,730–126,743)
IN	Black	137	22.8	4.32	(15.4–32.3)	||	42,152	(24,464–59,840)
IN	Other	134	13.3	3.27	(8.1–21.1)	||	24,285	(11,960–36,610)
KS	White	2,011	11.2	0.85	(9.7–13.0)	||	60,593	(51,459–69,727)
KS	Black	125	18.4	4.23	(11.5–28.1)	||	9,168	(4,683–13,653)
KS	Other	214	5.9	1.49	(3.6–9.6)	||	5,300	(2,679–7,921)
KY	White	1,605	8.5	0.84	(7.0–10.3)	||	64,710	(52,619–76,802)

Notes:
* CI denotes confidence interval.
** Total includes data from 31 states and DC, but excludes Puerto Rico.
If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.
When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

Asthma and Community Health Branch, National Center for Environmental Health
Centers for Disease Control and Prevention

Table L4
Child Lifetime Asthma Prevalence
and Weighted Number by Race and State or Territory: BRFSS 2018

State	Race	Unweighted Number	Prevalence (Percent)	Standard Error	95% CI * (Percent)		Weighted Number	95% CI* Weighted Number
KY	Black	126	22.5	6.40	(12.4–37.3)	||	21,614	(8,758–34,469)
KY	Other	133	4.5	1.73	(2.1–9.4)	||	5,210	(1,456–8,964)
ME	White	1,536	10.7	1.09	(8.7–13.0)	||	22,116	(17,642–26,591)
ME	Black	33	38.1	12.86	(17.5–64.2)	||	2,663	(274–5,052)
ME	Other	125	14.4	4.67	(7.4–26.1)	||	2,877	(883–4,871)
MD	White	630	8.9	1.51	(6.3–12.3)	||	45,443	(29,925–60,961)
MD	Black	260	15.5	2.87	(10.7–22.0)	||	68,856	(42,354–95,358)
MD	Other	166	17.9	3.73	(11.7–26.4)	||	47,781	(26,729–68,833)
MA	White	450	13.3	1.95	(9.9–17.6)	||	113,601	(80,382–146,821)
MA	Black	47	16.6	6.43	(7.4–33.1)	||	24,633	(4,889–44,377)
MA	Other	96	10.2	3.07	(5.6–18.0)	||	25,130	(10,190–40,071)
MI	White	1,761	11.4	0.95	(9.7–13.4)	||	161,358	(134,611–188,106)
MI	Black	319	18.1	2.64	(13.4–23.8)	||	65,664	(45,563–85,765)
MI	Other	301	10.8	2.04	(7.4–15.5)	||	31,323	(19,488–43,158)
MN	White	3,253	7.5	0.59	(6.4–8.7)	||	67,765	(57,077–78,453)
MN	Black	226	8.9	2.09	(5.5–13.9)	||	11,378	(5,953–16,803)
MN	Other	580	6.3	1.17	(4.4–9.1)	||	14,401	(9,146–19,657)
MS	White	718	6.8	0.99	(5.1–9.0)	||	21,869	(15,629–28,108)
MS	Black	633	14.2	1.63	(11.3–17.7)	||	43,275	(33,205–53,345)
MS	Other	64	4.4			||	1,628	
MO	White	813	10.3	1.39	(7.8–13.3)	||	98,012	(71,433–124,592)
MO	Black	111	23.1	5.31	(14.3–35.0)	||	46,303	(22,645–69,961)
MO	Other	69	22.2	7.06	(11.3–38.8)	||	30,287	(8,073–52,501)
MT	White	773	6.7	1.13	(4.8–9.3)	||	11,718	(7,804–15,632)
MT	Black	16	6.7			||	198	
MT	Other	365	9.0	2.45	(5.2–15.1)	||	3,826	(1,791–5,861)
NE	White	2,903	7.3	0.68	(6.1–8.8)	||	26,920	(22,019–31,822)
NE	Black	101	17.5	4.93	(9.8–29.3)	||	5,282	(2,072–8,493)
NE	Other	438	5.8	1.71	(3.2–10.2)	||	4,121	(1,667–6,575)
NV	White	488	10.9	2.76	(6.6–17.6)	||	36,260	(17,126–55,393)
NV	Black	39	17.9	7.10	(7.8–36.0)	||	15,500	(2,600–28,400)

Notes:
* CI denotes confidence interval.
** Total includes data from 31 states and DC, but excludes Puerto Rico.
If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.
When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

Asthma and Community Health Branch, National Center for Environmental Health
Centers for Disease Control and Prevention

Table L4
Child Lifetime Asthma Prevalence
and Weighted Number by Race and State or Territory: BRFSS 2018

State	Race	Unweighted Number	Prevalence (Percent)	Standard Error	95% CI * (Percent)	||	Weighted Number	95% CI* Weighted Number
NV	Other	205	14.8	3.42	(9.3–22.9)	||	35,921	(18,773–53,068)
NH	White	879	11.6	1.26	(9.3–14.3)	||	22,750	(17,873–27,627)
NH	Black	26	10.9			||	894	
NH	Other	75	8.7	4.04	(3.4–20.5)	||	2,164	(108–4,220)
NJ	White	237	11.4	2.76	(7.0–18.0)	||	118,960	(60,447–177,474)
NJ	Black	66	19.7	6.96	(9.3–36.7)	||	64,962	(21,402–108,522)
NJ	Other	83	5.5			||	26,888	
NM	White	1,101	11.7	1.45	(9.1–14.8)	||	39,752	(29,738–49,766)
NM	Black	48	19.1	9.24	(6.8–43.2)	||	3,624	(0–7,521)
NM	Other	424	10.1	1.85	(7.0–14.4)	||	11,073	(7,024–15,122)
NY	White	1,737	9.8	1.23	(7.6–12.5)	||	172,119	(128,397–215,840)
NY	Black	258	12.1	2.50	(8.0–17.9)	||	79,345	(46,558–112,132)
NY	Other	430	8.4	1.71	(5.6–12.4)	||	81,589	(48,538–114,641)
OH	White	2,110	9.8	0.90	(8.2–11.7)	||	173,714	(142,011–205,416)
OH	Black	212	20.7	3.58	(14.5–28.6)	||	84,537	(52,777–116,298)
OH	Other	194	14.5	3.62	(8.7–23.1)	||	39,033	(18,460–59,606)
OR	White	991	11.1	1.52	(8.5–14.5)	||	66,790	(48,231–85,349)
OR	Black	26	9.2			||	1,622	
OR	Other	219	13.0	3.68	(7.3–22.1)	||	25,758	(10,168–41,348)
PA	White	966	10.5	1.55	(7.8–14.0)	||	185,016	(129,707–240,325)
PA	Black	182	21.6	4.62	(13.9–32.0)	||	85,396	(44,777–126,016)
PA	Other	128	15.4	3.76	(9.4–24.3)	||	51,754	(26,112–77,396)
RI	White	788	14.1	2.28	(10.2–19.2)	||	17,907	(11,786–24,027)
RI	Black	81	9.9	3.36	(5.0–18.7)	||	2,037	(664–3,410)
RI	Other	168	16.9	3.86	(10.6–25.8)	||	6,829	(3,460–10,198)
UT	White	3,268	7.9	0.56	(6.9–9.1)	||	60,803	(52,211–69,395)
UT	Black	49	20.7	7.20	(10.0–38.2)	||	3,030	(710–5,349)
UT	Other	382	8.5	1.67	(5.7–12.4)	||	10,302	(6,195–14,409)
VT	White	1,121	12.7	1.36	(10.2–15.6)	||	11,202	(8,807–13,597)
VT	Black	12	21.4			||	302	
VT	Other	92	14.8	5.52	(6.9–29.1)	||	1,519	(324–2,714)

Notes:
* CI denotes confidence interval.
** Total includes data from 31 states and DC, but excludes Puerto Rico.
If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.
When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

Asthma and Community Health Branch, National Center for Environmental Health
Centers for Disease Control and Prevention

Table L4
Child Lifetime Asthma Prevalence
and Weighted Number by Race and State or Territory: BRFSS 2018

State	Race	Unweighted Number	Prevalence (Percent)	Standard Error	95% CI * (Percent)	‖	Weighted Number	95% CI* Weighted Number
WI	White	780	9.8	1.51	(7.2–13.2)	‖	90,360	(62,615–118,105)
WI	Black	50	14.6	5.76	(6.5–29.7)	‖	15,114	(2,895–27,333)
WI	Other	149	17.3	4.29	(10.5–27.4)	‖	34,930	(17,211–52,650)
Territory								

Notes:
* CI denotes confidence interval.
** Total includes data from 31 states and DC, but excludes Puerto Rico.
If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.
When the sample size is less than 50, estimates are not precise and should be interpreted with caution.