

Table L3
Adult* Lifetime Asthma Prevalence
and Number by Age and State or Territory: BRFSS 2019

State	Age Group	Sample Size [§]	Prevalence (percent)	Standard Error	95% CI [¶] (percent)	 	Weighted Number	95% CI [¶] (Weighted Number)
U.S. Total**	18-24	24,284	19.1	0.42	(18.2–19.9)		5,783,427	(5,509,853–6,057,000)
U.S. Total**	25-34	42,558	16.1	0.30	(15.5–16.7)		6,996,389	(6,720,067–7,272,711)
U.S. Total**	35-44	47,608	13.8	0.27	(13.3–14.3)		5,505,378	(5,285,557–5,725,198)
U.S. Total**	45-54	56,248	13.7	0.26	(13.2–14.2)		5,198,789	(4,991,362–5,406,215)
U.S. Total**	55-64	81,566	14.1	0.26	(13.6–14.6)		5,752,413	(5,529,776–5,975,051)
U.S. Total**	65+	149,544	12.2	0.18	(11.9–12.6)		6,424,889	(6,229,204–6,620,574)
AL	18-24	340	20.5	2.75	(15.0–25.9)		95,708	(66,908–124,507)
AL	25-34	678	18.5	1.79	(15.0–22.0)		117,024	(92,497–141,552)
AL	35-44	787	13.3	1.39	(10.6–16.1)		78,531	(61,451–95,611)
AL	45-54	1,015	13.6	1.31	(11.0–16.2)		80,630	(64,354–96,906)
AL	55-64	1,482	14.0	1.10	(11.8–16.1)		89,072	(74,514–103,629)
AL	65+	2,651	13.7	0.91	(12.0–15.5)		116,938	(100,763–133,113)
AK	18-24	168	13.7	3.22	(7.4–20.1)		9,749	(4,984–14,513)
AK	25-34	316	18.0	3.21	(11.7–24.3)		20,455	(12,516–28,395)
AK	35-44	414	13.5	2.29	(8.9–18.0)		12,581	(8,153–17,009)
AK	45-54	410	12.7	2.43	(7.9–17.5)		10,419	(6,225–14,612)
AK	55-64	697	14.2	1.96	(10.4–18.1)		12,999	(9,248–16,750)
AK	65+	903	14.6	1.75	(11.1–18.0)		13,396	(10,083–16,710)
AZ	18-24	482	18.1	2.38	(13.4–22.8)		127,136	(91,512–162,761)
AZ	25-34	794	17.9	1.90	(14.1–21.6)		174,658	(135,234–214,081)
AZ	35-44	903	14.2	1.56	(11.2–17.3)		125,880	(97,864–153,895)
AZ	45-54	1,120	16.1	1.69	(12.8–19.4)		134,259	(104,746–163,773)
AZ	55-64	1,686	17.1	1.52	(14.1–20.0)		146,523	(119,110–173,937)
AZ	65+	3,760	11.1	0.77	(9.6–12.6)		143,283	(123,339–163,227)
AR	18-24	201	21.2	3.29	(14.8–27.7)		61,463	(40,249–82,678)
AR	25-34	339	15.5	2.19	(11.1–19.8)		60,395	(42,326–78,463)
AR	35-44	484	13.2	1.69	(9.9–16.6)		48,706	(35,764–61,648)
AR	45-54	599	16.0	1.79	(12.4–19.5)		55,426	(42,177–68,675)
AR	55-64	1,007	13.4	1.27	(10.9–15.9)		50,208	(40,272–60,143)

Notes:

*Aged 18+ years

**U.S. Total includes 49 states plus the District of Columbia and excludes the three territories, New Jersey was excluded because of not meeting the minimum data collection requirements.

¶CI denotes confidence interval.

§Sample size is the number of respondents answered current asthma question

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the Sample Size is less than 50, estimates are not precise and should be interpreted with caution.

Table L3
Adult* Lifetime Asthma Prevalence
and Number by Age and State or Territory: BRFSS 2019

State	Age Group	Sample Size [§]	Prevalence (percent)	Standard Error	95% CI [¶] (percent)	 	Weighted Number	95% CI [¶] (Weighted Number)
AR	65+	2,622	12.3	0.83	(10.7–14.0)		64,277	(55,425–73,128)
CA	18-24	1,103	18.5	1.56	(15.5–21.6)		692,829	(564,780–820,877)
CA	25-34	1,860	15.0	0.96	(13.1–16.9)		885,354	(768,388–1,002,320)
CA	35-44	1,729	13.0	0.97	(11.1–14.9)		670,757	(567,626–773,888)
CA	45-54	1,799	12.0	0.89	(10.3–13.8)		596,560	(505,863–687,258)
CA	55-64	2,030	15.0	1.05	(12.9–17.0)		718,784	(612,815–824,753)
CA	65+	2,953	14.7	0.86	(13.0–16.4)		861,828	(756,731–966,925)
CO	18-24	611	21.8	1.85	(18.1–25.4)		116,906	(95,229–138,582)
CO	25-34	1,076	17.8	1.28	(15.3–20.3)		152,561	(129,010–176,113)
CO	35-44	1,095	14.5	1.18	(12.2–16.8)		113,485	(94,194–132,776)
CO	45-54	1,376	13.2	1.02	(11.2–15.2)		92,111	(77,335–106,887)
CO	55-64	1,769	13.2	0.90	(11.4–15.0)		94,838	(81,330–108,346)
CO	65+	3,202	13.1	0.69	(11.7–14.4)		110,930	(98,827–123,033)
CT	18-24	407	23.7	2.52	(18.8–28.7)		83,220	(63,362–103,079)
CT	25-34	710	22.3	1.92	(18.5–26.0)		97,485	(78,634–116,336)
CT	35-44	816	17.9	1.58	(14.8–21.0)		75,152	(60,993–89,311)
CT	45-54	1,357	15.3	1.23	(12.9–17.7)		68,458	(56,770–80,147)
CT	55-64	2,073	14.4	0.98	(12.5–16.3)		74,124	(63,580–84,669)
CT	65+	3,570	11.1	0.71	(9.7–12.5)		68,549	(59,562–77,536)
DE	18-24	247	23.2	3.91	(15.5–30.9)		20,343	(12,533–28,153)
DE	25-34	432	19.1	2.90	(13.4–24.8)		24,239	(16,074–32,404)
DE	35-44	477	14.1	2.21	(9.7–18.4)		15,831	(10,573–21,088)
DE	45-54	587	13.5	1.74	(10.0–16.9)		15,321	(11,234–19,408)
DE	55-64	774	16.5	1.76	(13.0–19.9)		21,941	(16,957–26,926)
DE	65+	1,302	11.1	1.14	(8.9–13.3)		20,748	(16,393–25,104)
DC	18-24	132	13.3	3.23	(7.0–19.7)		8,936	(4,365–13,506)
DC	25-34	350	23.1	2.51	(18.2–28.0)		37,196	(28,196–46,196)
DC	35-44	335	15.7	2.07	(11.7–19.8)		16,965	(12,309–21,620)
DC	45-54	362	17.3	2.22	(12.9–21.6)		12,439	(9,015–15,863)

Notes:

*Aged 18+ years

**U.S. Total includes 49 states plus the District of Columbia and excludes the three territories, New Jersey was excluded because of not meeting the minimum data collection requirements.

¶CI denotes confidence interval.

§Sample size is the number of respondents answered current asthma question

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the Sample Size is less than 50, estimates are not precise and should be interpreted with caution.

Table L3
Adult* Lifetime Asthma Prevalence
and Number by Age and State or Territory: BRFSS 2019

State	Age Group	Sample Size [§]	Prevalence (percent)	Standard Error	95% CI [¶] (percent)	 	Weighted Number	95% CI [¶] (Weighted Number)
DC	55-64	452	20.7	2.51	(15.8–25.7)		14,919	(10,864–18,975)
DC	65+	905	14.7	1.52	(11.7–17.7)		13,000	(10,159–15,840)
FL	18-24	932	13.1	1.71	(9.7–16.4)		235,104	(172,323–297,885)
FL	25-34	1,717	16.0	1.73	(12.6–19.4)		440,364	(339,436–541,291)
FL	35-44	1,788	13.2	1.39	(10.4–15.9)		334,127	(261,648–406,606)
FL	45-54	2,149	12.6	1.39	(9.9–15.3)		330,399	(256,272–404,526)
FL	55-64	3,122	13.1	1.50	(10.2–16.1)		373,094	(282,298–463,889)
FL	65+	6,899	10.5	0.75	(9.1–12.0)		477,595	(408,207–546,983)
GA	18-24	523	16.1	2.08	(12.0–20.2)		170,056	(124,050–216,063)
GA	25-34	823	13.2	1.64	(10.0–16.4)		190,670	(141,872–239,468)
GA	35-44	931	11.3	1.40	(8.5–14.0)		152,534	(113,854–191,215)
GA	45-54	1,071	11.0	1.26	(8.6–13.5)		149,203	(114,906–183,499)
GA	55-64	1,410	12.7	1.59	(9.5–15.8)		163,874	(120,078–207,669)
GA	65+	2,456	13.0	1.09	(10.9–15.2)		198,868	(164,183–233,553)
HI	18-24	481	21.5	2.17	(17.2–25.8)		25,944	(20,179–31,710)
HI	25-34	875	19.2	1.72	(15.8–22.5)		37,477	(30,138–44,817)
HI	35-44	1,020	15.8	1.43	(13.0–18.6)		29,201	(23,672–34,730)
HI	45-54	1,053	14.2	1.27	(11.7–16.7)		22,847	(18,614–27,079)
HI	55-64	1,473	14.3	1.17	(12.0–16.6)		25,300	(21,026–29,574)
HI	65+	2,686	15.0	1.02	(13.0–17.0)		39,535	(33,863–45,208)
ID	18-24	355	18.9	2.82	(13.3–24.4)		32,556	(22,016–43,095)
ID	25-34	609	16.7	2.13	(12.5–20.9)		37,870	(27,674–48,066)
ID	35-44	742	12.5	1.67	(9.2–15.7)		27,877	(20,198–35,555)
ID	45-54	719	15.0	2.05	(11.0–19.0)		27,753	(19,730–35,776)
ID	55-64	994	13.3	1.60	(10.2–16.5)		28,579	(21,439–35,719)
ID	65+	1,920	12.8	1.30	(10.2–15.3)		36,560	(28,713–44,407)
IL	18-24	443	17.5	2.37	(12.9–22.2)		210,284	(147,888–272,680)
IL	25-34	697	14.6	1.60	(11.4–17.7)		248,409	(190,595–306,223)
IL	35-44	808	15.0	1.45	(12.1–17.8)		243,551	(193,485–293,616)

Notes:

*Aged 18+ years

**U.S. Total includes 49 states plus the District of Columbia and excludes the three territories, New Jersey was excluded because of not meeting the minimum data collection requirements.

†CI denotes confidence interval.

§Sample size is the number of respondents answered current asthma question

¶If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the Sample Size is less than 50, estimates are not precise and should be interpreted with caution.

Table L3
Adult* Lifetime Asthma Prevalence
and Number by Age and State or Territory: BRFSS 2019

State	Age Group	Sample Size [§]	Prevalence (percent)	Standard Error	95% CI [¶] (percent)	 	Weighted Number	95% CI [¶] (Weighted Number)
IL	45-54	861	12.7	1.31	(10.1–15.3)		197,645	(155,200–240,090)
IL	55-64	1,049	13.9	1.17	(11.6–16.2)		227,770	(187,701–267,839)
IL	65+	1,595	10.1	0.85	(8.5–11.8)		205,999	(170,813–241,185)
IN	18-24	460	18.1	2.05	(14.1–22.2)		123,555	(93,088–154,022)
IN	25-34	799	15.2	1.35	(12.5–17.8)		130,036	(105,908–154,165)
IN	35-44	967	14.8	1.24	(12.4–17.3)		121,102	(99,830–142,374)
IN	45-54	1,235	14.5	1.13	(12.3–16.7)		110,256	(92,254–128,259)
IN	55-64	1,814	13.9	0.92	(12.1–15.7)		119,785	(103,219–136,351)
IN	65+	3,389	12.5	0.69	(11.1–13.8)		134,405	(118,913–149,896)
IA	18-24	710	17.6	1.61	(14.5–20.8)		57,887	(46,459–69,314)
IA	25-34	1,089	14.9	1.23	(12.5–17.3)		57,741	(47,582–67,901)
IA	35-44	1,235	12.6	1.04	(10.6–14.7)		47,872	(39,676–56,067)
IA	45-54	1,361	10.6	0.91	(8.8–12.4)		36,926	(30,429–43,423)
IA	55-64	1,909	10.7	0.79	(9.2–12.3)		43,992	(37,321–50,663)
IA	65+	3,331	8.7	0.56	(7.6–9.8)		47,812	(41,624–53,999)
KS	18-24	757	20.4	1.69	(17.1–23.7)		61,933	(50,830–73,035)
KS	25-34	1,222	16.6	1.19	(14.3–19.0)		61,642	(52,247–71,037)
KS	35-44	1,411	16.4	1.12	(14.2–18.6)		58,075	(49,666–66,484)
KS	45-54	1,509	14.2	1.07	(12.1–16.3)		44,297	(37,289–51,304)
KS	55-64	2,314	12.7	0.79	(11.2–14.3)		46,228	(40,292–52,163)
KS	65+	3,918	11.3	0.60	(10.1–12.5)		52,839	(47,143–58,534)
KY	18-24	502	19.1	2.22	(14.8–23.5)		82,175	(61,831–102,519)
KY	25-34	826	16.3	1.78	(12.8–19.8)		93,449	(72,110–114,788)
KY	35-44	888	10.4	1.35	(7.8–13.1)		56,710	(41,838–71,583)
KY	45-54	1,188	16.1	1.91	(12.3–19.8)		86,501	(64,158–108,844)
KY	55-64	1,664	14.4	1.23	(12.0–16.8)		84,050	(69,369–98,731)
KY	65+	2,774	11.3	0.92	(9.5–13.1)		85,905	(71,524–100,287)
LA	18-24	328	20.1	2.42	(15.4–24.9)		87,971	(65,679–110,263)
LA	25-34	585	17.1	1.72	(13.7–20.5)		111,545	(88,144–134,946)

Notes:

*Aged 18+ years

**U.S. Total includes 49 states plus the District of Columbia and excludes the three territories, New Jersey was excluded because of not meeting the minimum data collection requirements.

†CI denotes confidence interval.

§Sample size is the number of respondents answered current asthma question

¶If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the Sample Size is less than 50, estimates are not precise and should be interpreted with caution.

Table L3
Adult* Lifetime Asthma Prevalence
and Number by Age and State or Territory: BRFSS 2019

State	Age Group	Sample Size [§]	Prevalence (percent)	Standard Error	95% CI [¶] (percent)	 	Weighted Number	95% CI [¶] (Weighted Number)
LA	35-44	629	13.9	1.66	(10.6–17.2)		81,471	(61,085–101,857)
LA	45-54	749	13.9	1.44	(11.1–16.7)		72,139	(56,513–87,765)
LA	55-64	1,026	15.0	1.32	(12.4–17.6)		87,600	(71,398–103,801)
LA	65+	1,332	11.6	1.09	(9.4–13.7)		85,652	(69,054–102,250)
ME	18-24	359	23.8	3.43	(17.0–30.5)		26,807	(17,806–35,808)
ME	25-34	708	22.6	2.27	(18.1–27.0)		34,495	(26,662–42,329)
ME	35-44	926	18.8	1.90	(15.1–22.5)		28,740	(22,400–35,079)
ME	45-54	1,351	13.7	1.17	(11.4–16.0)		22,477	(18,563–26,391)
ME	55-64	2,512	12.6	0.88	(10.8–14.3)		26,692	(22,850–30,533)
ME	65+	5,270	13.0	0.66	(11.7–14.3)		35,934	(32,216–39,652)
MD	18-24	648	20.6	1.92	(16.8–24.4)		113,287	(90,299–136,276)
MD	25-34	1,316	16.9	1.26	(14.5–19.4)		136,080	(114,744–157,417)
MD	35-44	1,640	13.9	1.03	(11.9–15.9)		107,005	(90,622–123,388)
MD	45-54	2,376	14.9	0.97	(13.0–16.8)		112,269	(96,740–127,798)
MD	55-64	3,814	12.6	0.74	(11.1–14.0)		101,972	(89,454–114,490)
MD	65+	7,167	11.4	0.54	(10.4–12.5)		108,380	(98,022–118,738)
MA	18-24	557	21.1	2.05	(17.1–25.1)		149,414	(117,502–181,327)
MA	25-34	889	17.4	1.55	(14.4–20.5)		159,551	(129,009–190,092)
MA	35-44	904	14.3	1.29	(11.8–16.8)		118,323	(96,236–140,409)
MA	45-54	1,069	14.9	1.35	(12.2–17.5)		124,191	(100,362–148,021)
MA	55-64	1,524	14.1	1.08	(11.9–16.2)		132,553	(111,324–153,781)
MA	65+	2,510	13.2	0.86	(11.5–14.8)		152,923	(132,215–173,631)
MI	18-24	738	22.3	1.99	(18.4–26.2)		217,052	(173,467–260,637)
MI	25-34	1,094	17.3	1.41	(14.5–20.0)		219,179	(181,238–257,120)
MI	35-44	1,137	17.7	1.46	(14.8–20.5)		204,577	(167,898–241,255)
MI	45-54	1,454	16.2	1.20	(13.9–18.6)		192,503	(162,366–222,640)
MI	55-64	2,042	16.0	1.09	(13.9–18.2)		222,792	(190,191–255,393)
MI	65+	3,870	11.9	0.65	(10.6–13.2)		209,290	(185,848–232,732)
MN	18-24	1,160	16.0	1.22	(13.6–18.4)		83,115	(69,743–96,486)

Notes:

*Aged 18+ years

**U.S. Total includes 49 states plus the District of Columbia and excludes the three territories, New Jersey was excluded because of not meeting the minimum data collection requirements.

†CI denotes confidence interval.

§Sample size is the number of respondents answered current asthma question

¶If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the Sample Size is less than 50, estimates are not precise and should be interpreted with caution.

Table L3
Adult* Lifetime Asthma Prevalence
and Number by Age and State or Territory: BRFSS 2019

State	Age Group	Sample Size [§]	Prevalence (percent)	Standard Error	95% CI [¶] (percent)	 	Weighted Number	95% CI [¶] (Weighted Number)
MN	25-34	1,883	13.8	0.89	(12.0–15.5)		101,231	(87,546–114,915)
MN	35-44	2,254	13.9	0.82	(12.3–15.5)		99,586	(87,326–111,846)
MN	45-54	2,247	12.4	0.79	(10.9–14.0)		80,222	(69,673–90,771)
MN	55-64	3,077	12.5	0.72	(11.1–13.9)		94,218	(82,927–105,508)
MN	65+	4,663	10.5	0.53	(9.5–11.6)		96,840	(86,898–106,783)
MS	18-24	280	19.2	2.88	(13.6–24.9)		57,401	(38,396–76,405)
MS	25-34	540	12.1	1.65	(8.9–15.4)		46,969	(33,801–60,138)
MS	35-44	641	12.9	1.66	(9.7–16.2)		47,150	(34,473–59,828)
MS	45-54	738	16.1	1.80	(12.6–19.6)		55,112	(41,798–68,426)
MS	55-64	1,041	14.1	1.32	(11.5–16.7)		52,675	(42,439–62,911)
MS	65+	1,821	12.9	1.00	(11.0–14.9)		62,934	(52,916–72,953)
MO	18-24	444	22.0	2.40	(17.3–26.7)		128,017	(97,222–158,812)
MO	25-34	782	19.4	1.86	(15.8–23.1)		155,641	(123,197–188,085)
MO	35-44	738	14.0	1.52	(11.0–17.0)		104,634	(80,988–128,280)
MO	45-54	884	17.1	1.71	(13.7–20.4)		119,317	(93,619–145,016)
MO	55-64	1,378	12.0	1.08	(9.9–14.1)		97,477	(79,436–115,518)
MO	65+	2,936	10.1	0.71	(8.7–11.5)		106,599	(91,432–121,767)
MT	18-24	415	18.2	2.07	(14.1–22.3)		18,091	(13,647–22,534)
MT	25-34	657	19.4	1.71	(16.0–22.8)		26,239	(21,258–31,221)
MT	35-44	779	13.3	1.32	(10.7–15.8)		17,027	(13,503–20,551)
MT	45-54	752	14.7	1.48	(11.8–17.6)		16,832	(13,238–20,425)
MT	55-64	1,361	14.5	1.12	(12.3–16.7)		21,805	(18,280–25,331)
MT	65+	2,453	11.3	0.75	(9.8–12.7)		23,199	(20,052–26,346)
NE	18-24	965	15.5	1.45	(12.6–18.3)		30,304	(24,307–36,302)
NE	25-34	1,660	14.5	1.19	(12.2–16.8)		36,337	(30,036–42,639)
NE	35-44	1,918	13.2	1.04	(11.2–15.3)		31,918	(26,636–37,201)
NE	45-54	2,058	10.4	0.85	(8.7–12.1)		22,197	(18,477–25,916)
NE	55-64	3,093	12.1	0.84	(10.5–13.8)		28,993	(24,823–33,163)
NE	65+	6,035	10.1	0.55	(9.0–11.1)		31,540	(28,028–35,051)

Notes:

*Aged 18+ years

**U.S. Total includes 49 states plus the District of Columbia and excludes the three territories, New Jersey was excluded because of not meeting the minimum data collection requirements.

†CI denotes confidence interval.

§Sample size is the number of respondents answered current asthma question

¶If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the Sample Size is less than 50, estimates are not precise and should be interpreted with caution.

Table L3
Adult* Lifetime Asthma Prevalence
and Number by Age and State or Territory: BRFSS 2019

State	Age Group	Sample Size [§]	Prevalence (percent)	Standard Error	95% CI [¶] (percent)	 	Weighted Number	95% CI [¶] (Weighted Number)
NV	18-24	226	15.7	2.99	(9.8–21.6)		40,985	(24,443–57,527)
NV	25-34	387	17.3	2.37	(12.6–21.9)		74,513	(52,987–96,040)
NV	35-44	375	14.4	2.25	(10.0–18.8)		57,743	(38,935–76,551)
NV	45-54	378	11.7	2.08	(7.6–15.8)		44,156	(28,182–60,130)
NV	55-64	502	16.5	2.58	(11.5–21.6)		62,825	(41,535–84,116)
NV	65+	852	13.4	1.62	(10.2–16.6)		67,562	(50,712–84,412)
NH	18-24	185	22.8	3.50	(15.9–29.8)		25,752	(17,034–34,469)
NH	25-34	436	16.5	1.98	(12.6–20.3)		29,874	(22,433–37,315)
NH	35-44	500	22.5	2.35	(17.9–27.1)		34,639	(26,597–42,680)
NH	45-54	799	13.7	1.39	(11.0–16.4)		23,600	(18,673–28,526)
NH	55-64	1,299	13.9	1.12	(11.7–16.1)		29,707	(24,694–34,721)
NH	65+	2,501	12.0	0.81	(10.4–13.6)		30,094	(25,922–34,265)
NM	18-24	325	14.8	2.23	(10.4–19.2)		29,712	(20,322–39,101)
NM	25-34	585	17.1	1.91	(13.4–20.9)		47,083	(35,965–58,201)
NM	35-44	687	10.2	1.34	(7.5–12.8)		25,668	(18,784–32,551)
NM	45-54	778	11.2	1.38	(8.5–13.9)		26,021	(19,464–32,578)
NM	55-64	1,261	12.3	1.22	(9.9–14.7)		32,608	(25,914–39,302)
NM	65+	2,268	11.9	0.99	(9.9–13.8)		44,387	(36,756–52,018)
NY	18-24	835	22.2	1.83	(18.6–25.7)		396,899	(325,572–468,225)
NY	25-34	1,560	15.2	1.14	(13.0–17.5)		428,795	(361,111–496,479)
NY	35-44	1,576	13.3	1.06	(11.2–15.4)		319,303	(266,806–371,799)
NY	45-54	2,029	14.6	1.02	(12.6–16.6)		331,399	(283,101–379,697)
NY	55-64	2,819	13.9	0.90	(12.2–15.7)		351,280	(304,621–397,939)
NY	65+	4,964	12.3	0.71	(10.9–13.7)		394,374	(347,247–441,502)
NC	18-24	331	16.3	2.29	(11.8–20.8)		166,165	(116,031–216,299)
NC	25-34	599	13.4	1.50	(10.4–16.3)		183,839	(141,048–226,631)
NC	35-44	604	13.1	1.57	(10.0–16.1)		170,515	(127,619–213,410)
NC	45-54	701	13.1	1.40	(10.3–15.8)		166,455	(129,729–203,180)
NC	55-64	786	14.2	1.39	(11.5–16.9)		192,489	(153,447–231,531)

Notes:

*Aged 18+ years

**U.S. Total includes 49 states plus the District of Columbia and excludes the three territories, New Jersey was excluded because of not meeting the minimum data collection requirements.

†CI denotes confidence interval.

§Sample size is the number of respondents answered current asthma question

¶If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the Sample Size is less than 50, estimates are not precise and should be interpreted with caution.

Table L3
Adult* Lifetime Asthma Prevalence
and Number by Age and State or Territory: BRFSS 2019

State	Age Group	Sample Size [§]	Prevalence (percent)	Standard Error	95% CI [†] (percent)	 	Weighted Number	95% CI [†] (Weighted Number)
NC	65+	1,191	10.5	1.02	(8.5–12.5)		185,583	(149,047–222,118)
ND	18-24	278	16.3	2.59	(11.2–21.4)		13,489	(8,906–18,072)
ND	25-34	455	14.7	1.88	(11.0–18.4)		16,662	(12,189–21,135)
ND	35-44	541	15.2	1.98	(11.3–19.1)		14,171	(10,230–18,112)
ND	45-54	629	9.2	1.36	(6.5–11.8)		7,011	(4,893–9,130)
ND	55-64	1,225	9.4	1.08	(7.2–11.5)		8,707	(6,641–10,772)
ND	65+	2,332	9.9	0.77	(8.4–11.4)		11,846	(9,965–13,726)
OH	18-24	627	20.9	2.12	(16.7–25.1)		229,070	(178,397–279,743)
OH	25-34	1,086	17.5	1.47	(14.6–20.4)		263,080	(216,339–309,821)
OH	35-44	1,221	15.6	1.44	(12.8–18.4)		214,552	(172,769–256,334)
OH	45-54	1,788	16.7	1.35	(14.0–19.3)		232,129	(191,547–272,711)
OH	55-64	2,835	15.4	1.00	(13.4–17.4)		243,180	(209,954–276,406)
OH	65+	5,751	11.4	0.61	(10.2–12.6)		233,934	(208,253–259,615)
OK	18-24	326	21.2	2.51	(16.3–26.1)		82,588	(61,023–104,153)
OK	25-34	704	16.5	1.54	(13.5–19.5)		87,134	(69,970–104,297)
OK	35-44	828	15.4	1.50	(12.5–18.4)		75,782	(60,137–91,427)
OK	45-54	856	15.3	1.36	(12.6–18.0)		66,110	(53,787–78,432)
OK	55-64	1,115	15.6	1.29	(13.1–18.2)		75,142	(61,960–88,324)
OK	65+	2,383	13.2	0.85	(11.5–14.8)		83,154	(72,093–94,215)
OR	18-24	442	18.5	2.08	(14.4–22.6)		68,767	(52,340–85,193)
OR	25-34	812	19.9	1.62	(16.7–23.1)		115,201	(94,503–135,899)
OR	35-44	881	16.5	1.41	(13.7–19.2)		91,276	(74,633–107,918)
OR	45-54	905	15.6	1.40	(12.9–18.4)		75,461	(61,406–89,516)
OR	55-64	1,109	15.8	1.53	(12.7–18.8)		85,970	(67,862–104,078)
OR	65+	1,812	16.0	1.16	(13.7–18.3)		121,016	(102,473–139,560)
PA	18-24	485	25.0	2.55	(20.0–30.1)		299,655	(230,853–368,457)
PA	25-34	908	18.4	1.54	(15.4–21.4)		302,207	(248,383–356,032)
PA	35-44	816	17.7	1.68	(14.4–21.0)		266,360	(212,506–320,214)
PA	45-54	976	14.6	1.36	(11.9–17.2)		218,766	(176,270–261,262)

Notes:

*Aged 18+ years

**U.S. Total includes 49 states plus the District of Columbia and excludes the three territories, New Jersey was excluded because of not meeting the minimum data collection requirements.

†CI denotes confidence interval.

§Sample size is the number of respondents answered current asthma question

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the Sample Size is less than 50, estimates are not precise and should be interpreted with caution.

Table L3
Adult* Lifetime Asthma Prevalence
and Number by Age and State or Territory: BRFSS 2019

State	Age Group	Sample Size [§]	Prevalence (percent)	Standard Error	95% CI [¶] (percent)	 	Weighted Number	95% CI [¶] (Weighted Number)
PA	55-64	1,363	12.4	1.10	(10.3–14.6)		222,130	(181,333–262,927)
PA	65+	1,951	11.6	0.93	(9.8–13.4)		278,404	(232,581–324,226)
RI	18-24	257	20.3	3.20	(14.0–26.6)		21,858	(14,261–29,455)
RI	25-34	517	19.1	2.19	(14.8–23.4)		27,429	(20,663–34,195)
RI	35-44	573	14.4	1.83	(10.8–18.0)		17,917	(13,193–22,641)
RI	45-54	850	13.3	1.35	(10.6–16.0)		17,062	(13,518–20,606)
RI	55-64	1,388	17.1	1.29	(14.6–19.7)		25,457	(21,390–29,524)
RI	65+	2,402	14.2	0.89	(12.5–16.0)		26,445	(23,045–29,845)
SC	18-24	437	20.6	2.24	(16.2–25.0)		98,970	(75,520–122,421)
SC	25-34	658	16.8	1.80	(13.3–20.4)		112,863	(86,988–138,738)
SC	35-44	697	18.6	1.73	(15.2–22.0)		113,744	(91,038–136,449)
SC	45-54	906	12.9	1.41	(10.1–15.7)		76,930	(59,354–94,505)
SC	55-64	1,346	14.7	1.28	(12.2–17.2)		99,228	(80,929–117,527)
SC	65+	2,923	12.0	0.86	(10.3–13.7)		110,957	(94,496–127,418)
SD	18-24	343	24.2	4.18	(16.0–32.4)		20,843	(12,611–29,076)
SD	25-34	621	10.2	1.92	(6.4–14.0)		11,459	(7,063–15,854)
SD	35-44	749	10.8	1.92	(7.0–14.5)		11,411	(7,282–15,540)
SD	45-54	888	10.8	1.92	(7.0–14.6)		9,812	(6,239–13,384)
SD	55-64	1,316	12.3	1.68	(9.0–15.6)		14,029	(10,070–17,989)
SD	65+	2,594	7.7	0.99	(5.7–9.6)		11,737	(8,717–14,756)
TN	18-24	370	19.2	2.53	(14.2–24.1)		119,307	(85,670–152,943)
TN	25-34	677	17.3	1.78	(13.8–20.8)		159,338	(125,086–193,590)
TN	35-44	700	13.6	1.58	(10.5–16.7)		111,739	(84,893–138,584)
TN	45-54	1,021	16.0	1.48	(13.1–18.8)		130,919	(105,511–156,328)
TN	55-64	1,308	15.3	1.29	(12.8–17.9)		135,460	(111,531–159,389)
TN	65+	2,036	12.3	0.90	(10.6–14.1)		142,306	(121,153–163,459)
TX	18-24	797	17.9	2.12	(13.7–22.1)		519,159	(388,508–649,811)
TX	25-34	1,460	15.2	1.60	(12.1–18.4)		623,472	(484,512–762,432)
TX	35-44	1,524	9.4	1.18	(7.1–11.7)		362,172	(270,437–453,908)

Notes:

*Aged 18+ years

**U.S. Total includes 49 states plus the District of Columbia and excludes the three territories, New Jersey was excluded because of not meeting the minimum data collection requirements.

†CI denotes confidence interval.

§Sample size is the number of respondents answered current asthma question

¶If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the Sample Size is less than 50, estimates are not precise and should be interpreted with caution.

Table L3
Adult* Lifetime Asthma Prevalence
and Number by Age and State or Territory: BRFSS 2019

State	Age Group	Sample Size [§]	Prevalence (percent)	Standard Error	95% CI [¶] (percent)	 	Weighted Number	95% CI [¶] (Weighted Number)
TX	45-54	1,655	12.1	1.43	(9.3–14.9)		412,444	(311,597–513,291)
TX	55-64	2,168	13.2	1.58	(10.1–16.3)		433,342	(323,039–543,645)
TX	65+	4,478	12.9	1.08	(10.8–15.0)		494,506	(408,348–580,663)
UT	18-24	1,030	20.4	1.47	(17.5–23.3)		74,547	(62,795–86,299)
UT	25-34	1,668	16.7	1.07	(14.6–18.8)		76,735	(66,282–87,187)
UT	35-44	2,006	14.7	0.95	(12.9–16.6)		63,918	(55,301–72,535)
UT	45-54	1,715	13.5	1.01	(11.6–15.5)		43,646	(36,817–50,475)
UT	55-64	1,870	15.4	1.02	(13.4–17.4)		46,027	(39,563–52,491)
UT	65+	3,263	13.0	0.72	(11.6–14.4)		47,589	(42,147–53,032)
VT	18-24	225	21.4	3.19	(15.1–27.7)		14,227	(9,600–18,854)
VT	25-34	466	23.8	2.54	(18.8–28.8)		17,267	(13,291–21,243)
VT	35-44	633	21.2	2.14	(17.0–25.4)		14,906	(11,642–18,171)
VT	45-54	893	14.6	1.55	(11.6–17.7)		10,804	(8,377–13,230)
VT	55-64	1,494	17.4	1.40	(14.6–20.1)		16,594	(13,699–19,488)
VT	65+	2,611	13.1	0.96	(11.2–15.0)		16,198	(13,714–18,682)
VA	18-24	636	16.6	1.70	(13.3–20.0)		140,026	(109,933–170,120)
VA	25-34	1,041	15.4	1.37	(12.7–18.1)		174,721	(142,407–207,034)
VA	35-44	1,152	14.8	1.30	(12.2–17.3)		160,467	(130,906–190,027)
VA	45-54	1,421	12.9	1.13	(10.7–15.1)		137,386	(112,399–162,374)
VA	55-64	2,024	13.2	0.91	(11.5–15.0)		147,575	(126,744–168,407)
VA	65+	3,488	12.7	0.77	(11.2–14.2)		171,998	(150,232–193,764)
WA	18-24	716	21.8	1.81	(18.2–25.4)		146,894	(119,977–173,810)
WA	25-34	1,434	17.6	1.13	(15.4–19.8)		194,813	(168,412–221,214)
WA	35-44	1,619	15.1	1.01	(13.1–17.1)		150,357	(129,249–171,466)
WA	45-54	1,700	15.7	1.04	(13.7–17.8)		139,189	(119,714–158,663)
WA	55-64	2,508	16.9	0.99	(15.0–18.9)		164,641	(144,107–185,174)
WA	65+	4,721	14.3	0.68	(13.0–15.7)		175,250	(157,924–192,576)
WV	18-24	199	20.2	3.08	(14.1–26.3)		32,619	(21,777–43,461)
WV	25-34	383	18.7	2.23	(14.3–23.1)		38,804	(28,795–48,813)

Notes:

*Aged 18+ years

**U.S. Total includes 49 states plus the District of Columbia and excludes the three territories, New Jersey was excluded because of not meeting the minimum data collection requirements.

†CI denotes confidence interval.

§Sample size is the number of respondents answered current asthma question

¶If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the Sample Size is less than 50, estimates are not precise and should be interpreted with caution.

Table L3
Adult* Lifetime Asthma Prevalence
and Number by Age and State or Territory: BRFSS 2019

State	Age Group	Sample Size [§]	Prevalence (percent)	Standard Error	95% CI [¶] (percent)	 	Weighted Number	95% CI [¶] (Weighted Number)
WV	35-44	573	16.9	1.79	(13.4–20.4)		35,697	(27,658–43,736)
WV	45-54	794	18.8	1.72	(15.5–22.2)		41,388	(33,143–49,632)
WV	55-64	1,142	17.3	1.36	(14.6–19.9)		42,859	(35,650–50,069)
WV	65+	2,131	12.4	0.86	(10.7–14.0)		45,229	(38,822–51,635)
WI	18-24	277	18.1	2.67	(12.9–23.4)		102,760	(70,287–135,234)
WI	25-34	425	14.7	2.03	(10.7–18.7)		105,716	(74,989–136,443)
WI	35-44	546	16.5	1.95	(12.6–20.3)		115,853	(86,469–145,237)
WI	45-54	600	13.7	1.81	(10.1–17.2)		91,088	(65,607–116,569)
WI	55-64	1,134	13.6	1.28	(11.1–16.2)		111,230	(89,289–133,171)
WI	65+	1,873	11.5	0.99	(9.5–13.4)		115,366	(94,881–135,851)
WY	18-24	189	21.7	3.53	(14.7–28.7)		11,893	(7,543–16,242)
WY	25-34	350	14.9	2.07	(10.9–19.0)		11,090	(7,868–14,312)
WY	35-44	411	10.7	1.73	(7.3–14.1)		7,823	(5,225–10,420)
WY	45-54	517	13.8	1.89	(10.1–17.5)		8,305	(5,917–10,693)
WY	55-64	967	11.9	1.36	(9.2–14.5)		9,077	(6,923–11,230)
WY	65+	2,126	11.5	0.98	(9.6–13.5)		11,196	(9,235–13,157)
Territories								
GU	18-24	237	12.4	2.39	(7.7–17.1)		2,184	(1,342–3,026)
GU	25-34	388	7.2	1.49	(4.3–10.2)		1,524	(898–2,151)
GU	35-44	357	10.3	1.88	(6.6–14.0)		2,309	(1,469–3,148)
GU	45-54	444	13.8	4.13	(5.7–21.9)		2,718	(909–4,528)
GU	55-64	459	11.0	3.28	(4.6–17.4)		1,543	(557–2,530)
GU	65+	520	7.4	1.45	(4.6–10.3)		799	(503–1,095)
PR	18-24	446	18.7	2.10	(14.6–22.9)		64,912	(49,184–80,640)
PR	25-34	794	21.6	1.83	(18.0–25.1)		93,576	(76,554–110,598)
PR	35-44	895	19.5	1.62	(16.3–22.7)		85,450	(70,121–100,779)
PR	45-54	1,014	15.7	1.35	(13.1–18.3)		69,155	(56,648–81,663)
PR	55-64	1,090	19.4	1.48	(16.5–22.3)		83,675	(69,957–97,393)
PR	65+	1,750	15.3	1.07	(13.2–17.4)		95,945	(81,857–110,033)

Notes:

*Aged 18+ years

**U.S. Total includes 49 states plus the District of Columbia and excludes the three territories, New Jersey was excluded because of not meeting the minimum data collection requirements.

¶CI denotes confidence interval.

§Sample size is the number of respondents answered current asthma question

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the Sample Size is less than 50, estimates are not precise and should be interpreted with caution.