

Table C1
Child[§] Current Asthma Prevalence
and Weighted Number by State or Territory: BRFSS 2019

State	Sample Size	Prevalence (Percent)	Standard Error	95% CI [¶] (Percent)	 	Weighted Number	95% CI [¶] (Number)
Total *	47,807	7.4	0.26	(6.9–7.9)		3,622,537	(3,363,789–3,881,286)
CA	2,679	7.3	0.63	(6.2–8.7)		645,442	(535,011–755,874)
CT	1,493	8.8	0.95	(7.1–10.8)		61,005	(47,908–74,101)
DC	442	11.7	2.07	(8.2–16.4)		14,823	(9,492–20,155)
FL	2,576	6.7	0.86	(5.2–8.6)		269,934	(201,329–338,539)
GA	1,343	10.1	1.48	(7.6–13.4)		240,431	(169,434–311,429)
HI	1,688	7.5	0.76	(6.1–9.1)		21,411	(17,082–25,740)
IL	1,267	6.9	0.95	(5.3–9.0)		188,519	(136,914–240,124)
IN	1,650	6.7	0.68	(5.5–8.2)		99,608	(79,823–119,394)
KS	2,538	8.7	0.78	(7.3–10.4)		59,161	(48,550–69,771)
ME	1,494	9.4	1.15	(7.3–11.9)		21,291	(16,060–26,522)
MA	666	9.5	1.73	(6.6–13.5)		118,982	(74,800–163,163)
MI	2,127	7.8	0.78	(6.4–9.5)		160,490	(128,518–192,462)
MN	3,322	5.1	0.49	(4.2–6.2)		64,035	(52,043–76,027)
MS	1,090	7.6	1.10	(5.7–10.1)		50,586	(36,054–65,119)
MO	1,313	7.0	0.94	(5.4–9.1)		91,503	(67,394–115,611)
MT	1,423	6.7	0.82	(5.3–8.5)		15,016	(11,406–18,626)
NE	3,504	6.2	0.58	(5.2–7.4)		29,079	(23,749–34,409)
NV	665	7.9	1.71	(5.1–11.9)		50,434	(28,339–72,528)
NH	884	7.3	1.11	(5.4–9.7)		17,863	(12,493–23,232)
NM	1,226	5.4	1.02	(3.8–7.8)		25,705	(16,175–35,235)
NY	800	8.1	1.44	(5.7–11.4)		315,260	(202,022–428,498)
OH	2,160	7.7	0.89	(6.1–9.7)		190,815	(146,830–234,799)
OR	1,195	6.9	1.41	(4.6–10.3)		55,007	(32,284–77,730)
PA	1,375	8.0	0.84	(6.5–9.8)		198,040	(157,196–238,884)
RI	892	8.7	1.36	(6.3–11.7)		17,013	(11,653–22,374)
TX	2,377	6.8	0.98	(5.1–9.0)		488,562	(348,954–628,170)
UT	3,780	5.7	0.44	(4.9–6.7)		52,667	(44,782–60,551)
VT	1,064	8.1	1.12	(6.2–10.6)		9,131	(6,644–11,617)

Notes:

[§]Aged 0-17

[¶]CI denotes confidence interval.

* Total includes data from 29 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

**Asthma and Community Health Branch, National Center for Environmental Health
Centers for Disease Control and Prevention**

Table C1
Child[§] Current Asthma Prevalence
and Weighted Number by State or Territory: BRFSS 2019

State	Sample Size	Prevalence (Percent)	Standard Error	95% CI [*] (Percent)	 	Weighted Number	95% CI [*] (Number)
WI	774	4.3	0.97	(2.7–6.6)		50,725	(28,120–73,330)
Territory							
PR	1,418	15.0	1.08	(13.0–17.2)		103,946	(88,925–118,968)

Notes:

[§]Aged 0-17

^{*}CI denotes confidence interval.

* Total includes data from 29 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

**Asthma and Community Health Branch, National Center for Environmental Health
Centers for Disease Control and Prevention**