

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, all ages				
...	All causes	2,426,264	100.0	810.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	631,636	26.0	211.0
2	Malignant neoplasms (C00-C97)	559,888	23.1	187.0
3	Cerebrovascular diseases (I60-I69)	137,119	5.7	45.8
4	Chronic lower respiratory diseases (J40-J47)	124,583	5.1	41.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121,599	5.0	40.6
6	Diabetes mellitus (E10-E14)	72,449	3.0	24.2
7	Alzheimer's disease (G30)	72,432	3.0	24.2
8	Influenza and pneumonia (J10-J18)	56,326	2.3	18.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45,344	1.9	15.1
10	Septicemia (A40-A41)	34,234	1.4	11.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33,300	1.4	11.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	27,555	1.1	9.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	23,855	1.0	8.0
14	Parkinson's disease (G20-G21)	19,566	0.8	6.5
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18,573	0.8	6.2
...	All other causes (Residual)	447,805	18.5	149.6
All races, both sexes, 1-4 years				
...	All causes	4,631	100.0	28.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,610	34.8	9.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	515	11.1	3.2
3	Malignant neoplasms (C00-C97)	377	8.1	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	366	7.9	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	161	3.5	1.0
6	Influenza and pneumonia (J10-J18)	125	2.7	0.8
7	Septicemia (A40-A41)	88	1.9	0.5
8	Certain conditions originating in the perinatal period (P00-P96)	65	1.4	0.4
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	60	1.3	0.4
10	Cerebrovascular diseases (I60-I69)	54	1.2	0.3
11	Chronic lower respiratory diseases (J40-J47)	43	0.9	0.3
12	Anemias (D50-D64)	26	0.6	0.2
13	Complications of medical and surgical care (Y40-Y84,Y88)	22	0.5	0.1
14	Meningitis (G00,G03)	20	0.4	0.1
15	Meningococcal infection (A39)	14	0.3	*
15	Acute bronchitis and bronchiolitis (J20-J21)	14	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.3	*
...	All other causes (Residual)	1,057	22.8	6.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 5-14 years				
...	All causes	6,149	100.0	15.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,258	36.7	5.6
2	Malignant neoplasms (C00-C97)	907	14.8	2.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	390	6.3	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	344	5.6	0.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	253	4.1	0.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	219	3.6	0.5
7	Chronic lower respiratory diseases (J40-J47)	115	1.9	0.3
8	Cerebrovascular diseases (I60-I69)	95	1.5	0.2
9	Septicemia (A40-A41)	84	1.4	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	76	1.2	0.2
11	Influenza and pneumonia (J10-J18)	68	1.1	0.2
12	Anemias (D50-D64)	34	0.6	0.1
13	Diabetes mellitus (E10-E14)	28	0.5	0.1
14	Certain conditions originating in the perinatal period (P00-P96)	24	0.4	0.1
15	Complications of medical and surgical care (Y40-Y84,Y88)	19	0.3	*
...	All other causes (Residual)	1,235	20.1	3.1
All races, both sexes, 15-24 years				
...	All causes	34,887	100.0	82.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,229	46.5	38.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5,717	16.4	13.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,189	12.0	9.9
4	Malignant neoplasms (C00-C97)	1,644	4.7	3.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,076	3.1	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	460	1.3	1.1
7	Cerebrovascular diseases (I60-I69)	210	0.6	0.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	206	0.6	0.5
9	Influenza and pneumonia (J10-J18)	184	0.5	0.4
10	Pregnancy, childbirth and the puerperium (O00-099)	179	0.5	0.4
11	Diabetes mellitus (E10-E14)	165	0.5	0.4
12	Chronic lower respiratory diseases (J40-J47)	151	0.4	0.4
13	Septicemia (A40-A41)	139	0.4	0.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	100	0.3	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	99	0.3	0.2
...	All other causes (Residual)	4,139	11.9	9.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 25-34 years				
...	All causes	42,952	100.0	106.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,954	34.8	37.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,985	11.6	12.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,725	11.0	11.7
4	Malignant neoplasms (C00-C97)	3,656	8.5	9.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,307	7.7	8.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,182	2.8	2.9
7	Diabetes mellitus (E10-E14)	673	1.6	1.7
8	Cerebrovascular diseases (I60-I69)	527	1.2	1.3
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	437	1.0	1.1
10	Influenza and pneumonia (J10-J18)	335	0.8	0.8
11	Pregnancy, childbirth and the puerperium (O00-099)	324	0.8	0.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	316	0.7	0.8
13	Septicemia (A40-A41)	291	0.7	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	276	0.6	0.7
15	Chronic lower respiratory diseases (J40-J47)	255	0.6	0.6
...	All other causes (Residual)	6,709	15.6	16.6
All races, both sexes, 35-44 years				
...	All causes	83,043	100.0	190.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17,534	21.1	40.2
2	Malignant neoplasms (C00-C97)	13,917	16.8	31.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,339	14.9	28.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,591	7.9	15.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	4,010	4.8	9.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,020	3.6	6.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,551	3.1	5.8
8	Cerebrovascular diseases (I60-I69)	2,221	2.7	5.1
9	Diabetes mellitus (E10-E14)	2,094	2.5	4.8
10	Septicemia (A40-A41)	870	1.0	2.0
11	Influenza and pneumonia (J10-J18)	841	1.0	1.9
11	Chronic lower respiratory diseases (J40-J47)	841	1.0	1.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	785	0.9	1.8
14	Viral hepatitis (B15-B19)	534	0.6	1.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	468	0.6	1.1
...	All other causes (Residual)	14,427	17.4	33.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 45-54 years				
...	All causes	185,031	100.0	427.5
1	Malignant neoplasms (C00-C97)	50,334	27.2	116.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	38,095	20.6	88.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19,675	10.6	45.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	7,712	4.2	17.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,426	4.0	17.2
6	Cerebrovascular diseases (I60-I69)	6,341	3.4	14.7
7	Diabetes mellitus (E10-E14)	5,692	3.1	13.2
8	Human immunodeficiency virus (HIV) disease (B20-B24)	4,377	2.4	10.1
9	Chronic lower respiratory diseases (J40-J47)	3,924	2.1	9.1
10	Viral hepatitis (B15-B19)	2,911	1.6	6.7
11	Septicemia (A40-A41)	2,260	1.2	5.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,245	1.2	5.2
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,209	1.2	5.1
14	Influenza and pneumonia (J10-J18)	2,007	1.1	4.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,299	0.7	3.0
...	All other causes (Residual)	28,524	15.4	65.9
All races, both sexes, 55-64 years				
...	All causes	281,401	100.0	890.9
1	Malignant neoplasms (C00-C97)	101,454	36.1	321.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	65,477	23.3	207.3
3	Chronic lower respiratory diseases (J40-J47)	12,375	4.4	39.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,446	4.1	36.2
5	Diabetes mellitus (E10-E14)	11,432	4.1	36.2
6	Cerebrovascular diseases (I60-I69)	10,518	3.7	33.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	7,217	2.6	22.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,583	1.6	14.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,368	1.6	13.8
10	Septicemia (A40-A41)	4,032	1.4	12.8
11	Influenza and pneumonia (J10-J18)	3,154	1.1	10.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,178	0.8	6.9
13	Viral hepatitis (B15-B19)	2,127	0.8	6.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,722	0.6	5.5
15	Aortic aneurysm and dissection (I71)	1,366	0.5	4.3
...	All other causes (Residual)	37,952	13.5	120.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 65-74 years				
...	All causes	390,093	100.0	2,062.1
1	Malignant neoplasms (C00-C97)	137,554	35.3	727.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	92,752	23.8	490.3
3	Chronic lower respiratory diseases (J40-J47)	28,236	7.2	149.3
4	Cerebrovascular diseases (I60-I69)	18,223	4.7	96.3
5	Diabetes mellitus (E10-E14)	15,483	4.0	81.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,420	2.2	44.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,453	1.9	39.4
8	Septicemia (A40-A41)	6,064	1.6	32.1
9	Influenza and pneumonia (J10-J18)	6,061	1.6	32.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	4,917	1.3	26.0
11	Alzheimer's disease (G30)	3,812	1.0	20.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,174	0.8	16.8
13	Aortic aneurysm and dissection (I71)	2,635	0.7	13.9
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,384	0.6	12.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,369	0.6	12.5
...	All other causes (Residual)	50,556	13.0	267.3
All races, both sexes, 75-84 years				
...	All causes	667,338	100.0	5,115.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	180,451	27.0	1,383.1
2	Malignant neoplasms (C00-C97)	164,889	24.7	1,263.8
3	Chronic lower respiratory diseases (J40-J47)	47,406	7.1	363.4
4	Cerebrovascular diseases (I60-I69)	43,719	6.6	335.1
5	Alzheimer's disease (G30)	22,915	3.4	175.6
6	Diabetes mellitus (E10-E14)	21,763	3.3	166.8
7	Influenza and pneumonia (J10-J18)	16,668	2.5	127.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,536	2.2	111.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,708	2.1	105.1
10	Septicemia (A40-A41)	10,745	1.6	82.4
11	Parkinson's disease (G20-G21)	9,101	1.4	69.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6,654	1.0	51.0
13	Pneumonitis due to solids and liquids (J69)	5,284	0.8	40.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,750	0.7	36.4
15	Aortic aneurysm and dissection (I71)	4,622	0.7	35.4
...	All other causes (Residual)	100,127	15.0	767.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 85 years and over				
...	All causes	701,992	100.0	13,253.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	237,339	33.8	4,480.8
2	Malignant neoplasms (C00-C97)	85,072	12.1	1,606.1
3	Cerebrovascular diseases (I60-I69)	55,068	7.8	1,039.6
4	Alzheimer's disease (G30)	44,933	6.4	848.3
5	Chronic lower respiratory diseases (J40-J47)	31,203	4.4	589.1
6	Influenza and pneumonia (J10-J18)	26,617	3.8	502.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,388	2.2	290.5
8	Diabetes mellitus (E10-E14)	15,105	2.2	285.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,561	2.1	274.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10,030	1.4	189.4
11	Septicemia (A40-A41)	9,392	1.3	177.3
12	Pneumonitis due to solids and liquids (J69)	8,254	1.2	155.8
13	Parkinson's disease (G20-G21)	7,669	1.1	144.8
14	Atherosclerosis (I70)	4,764	0.7	89.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,134	0.6	78.0
...	All other causes (Residual)	132,463	18.9	2,500.8
All races, male, all ages				
...	All causes	1,201,942	100.0	814.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	315,706	26.3	214.0
2	Malignant neoplasms (C00-C97)	290,069	24.1	196.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	78,941	6.6	53.5
4	Chronic lower respiratory diseases (J40-J47)	59,260	4.9	40.2
5	Cerebrovascular diseases (I60-I69)	54,524	4.5	37.0
6	Diabetes mellitus (E10-E14)	36,006	3.0	24.4
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26,308	2.2	17.8
8	Influenza and pneumonia (J10-J18)	25,650	2.1	17.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22,094	1.8	15.0
10	Alzheimer's disease (G30)	21,151	1.8	14.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	17,866	1.5	12.1
12	Septicemia (A40-A41)	15,522	1.3	10.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14,717	1.2	10.0
14	Parkinson's disease (G20-G21)	11,300	0.9	7.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,415	0.8	6.4
...	All other causes (Residual)	203,413	16.9	137.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 1-4 years				
...	All causes	2,541	100.0	30.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	927	36.5	11.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	244	9.6	2.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	208	8.2	2.5
4	Malignant neoplasms (C00-C97)	207	8.1	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	89	3.5	1.1
6	Influenza and pneumonia (J10-J18)	63	2.5	0.8
7	Septicemia (A40-A41)	43	1.7	0.5
8	Certain conditions originating in the perinatal period (P00-P96)	37	1.5	0.4
9	Chronic lower respiratory diseases (J40-J47)	29	1.1	0.3
10	Cerebrovascular diseases (I60-I69)	26	1.0	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.9	0.3
12	Anemias (D50-D64)	15	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.4	*
14	Meningococcal infection (A39)	10	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.4	*
...	All other causes (Residual)	599	23.6	7.2
All races, male, 5-14 years				
...	All causes	3,625	100.0	17.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,385	38.2	6.7
2	Malignant neoplasms (C00-C97)	507	14.0	2.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	246	6.8	1.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	181	5.0	0.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	153	4.2	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	140	3.9	0.7
7	Chronic lower respiratory diseases (J40-J47)	78	2.2	0.4
8	Cerebrovascular diseases (I60-I69)	56	1.5	0.3
9	Septicemia (A40-A41)	52	1.4	0.3
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	1.1	0.2
11	Influenza and pneumonia (J10-J18)	35	1.0	0.2
12	Anemias (D50-D64)	16	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	14	0.4	*
14	Diabetes mellitus (E10-E14)	13	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.3	*
...	All other causes (Residual)	696	19.2	3.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 15-24 years				
...	All causes	26,070	100.0	119.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,234	46.9	56.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,991	19.1	22.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,528	13.5	16.2
4	Malignant neoplasms (C00-C97)	1,007	3.9	4.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	711	2.7	3.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	279	1.1	1.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	121	0.5	0.6
8	Cerebrovascular diseases (I60-I69)	115	0.4	0.5
9	Influenza and pneumonia (J10-J18)	101	0.4	0.5
10	Diabetes mellitus (E10-E14)	96	0.4	0.4
11	Chronic lower respiratory diseases (J40-J47)	93	0.4	0.4
12	Legal intervention (Y35,Y89.0)	82	0.3	0.4
13	Septicemia (A40-A41)	68	0.3	0.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	0.2	0.3
15	Anemias (D50-D64)	55	0.2	0.3
...	All other causes (Residual)	2,533	9.7	11.6
All races, male, 25-34 years				
...	All causes	30,188	100.0	146.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,466	38.0	55.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,053	13.4	19.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,983	13.2	19.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,298	7.6	11.2
5	Malignant neoplasms (C00-C97)	1,770	5.9	8.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	729	2.4	3.5
7	Diabetes mellitus (E10-E14)	376	1.2	1.8
8	Cerebrovascular diseases (I60-I69)	290	1.0	1.4
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	263	0.9	1.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	215	0.7	1.0
11	Influenza and pneumonia (J10-J18)	175	0.6	0.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	157	0.5	0.8
13	Septicemia (A40-A41)	152	0.5	0.7
14	Legal intervention (Y35,Y89.0)	139	0.5	0.7
15	Chronic lower respiratory diseases (J40-J47)	137	0.5	0.7
...	All other causes (Residual)	3,985	13.2	19.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 35-44 years				
...	All causes	52,150	100.0	238.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,431	23.8	56.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,625	16.5	39.5
3	Malignant neoplasms (C00-C97)	5,979	11.5	27.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,065	9.7	23.2
5	Human immunodeficiency virus (HIV) disease (B20-B24)	2,829	5.4	12.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,269	4.4	10.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,666	3.2	7.6
8	Diabetes mellitus (E10-E14)	1,279	2.5	5.9
9	Cerebrovascular diseases (I60-I69)	1,163	2.2	5.3
10	Influenza and pneumonia (J10-J18)	462	0.9	2.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	458	0.9	2.1
12	Septicemia (A40-A41)	452	0.9	2.1
13	Chronic lower respiratory diseases (J40-J47)	396	0.8	1.8
14	Viral hepatitis (B15-B19)	370	0.7	1.7
15	Aortic aneurysm and dissection (I71)	269	0.5	1.2
...	All other causes (Residual)	8,437	16.2	38.6
All races, male, 45-54 years				
...	All causes	115,172	100.0	541.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,433	23.8	128.9
2	Malignant neoplasms (C00-C97)	25,341	22.0	119.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,837	12.0	65.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,576	4.8	26.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,517	4.8	25.9
6	Diabetes mellitus (E10-E14)	3,515	3.1	16.5
7	Cerebrovascular diseases (I60-I69)	3,482	3.0	16.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	3,263	2.8	15.3
9	Viral hepatitis (B15-B19)	2,102	1.8	9.9
10	Chronic lower respiratory diseases (J40-J47)	1,958	1.7	9.2
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,628	1.4	7.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,338	1.2	6.3
13	Septicemia (A40-A41)	1,247	1.1	5.9
14	Influenza and pneumonia (J10-J18)	1,186	1.0	5.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	813	0.7	3.8
...	All other causes (Residual)	16,936	14.7	79.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 55-64 years				
...	All causes	168,987	100.0	1,110.0
1	Malignant neoplasms (C00-C97)	55,348	32.8	363.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	45,177	26.7	296.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,848	4.6	51.6
4	Diabetes mellitus (E10-E14)	6,670	3.9	43.8
5	Chronic lower respiratory diseases (J40-J47)	6,382	3.8	41.9
6	Cerebrovascular diseases (I60-I69)	5,897	3.5	38.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	5,095	3.0	33.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,463	2.0	22.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,442	1.4	16.0
10	Septicemia (A40-A41)	2,144	1.3	14.1
11	Influenza and pneumonia (J10-J18)	1,877	1.1	12.3
12	Viral hepatitis (B15-B19)	1,502	0.9	9.9
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,352	0.8	8.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,273	0.8	8.4
15	Aortic aneurysm and dissection (I71)	1,004	0.6	6.6
...	All other causes (Residual)	21,513	12.7	141.3
All races, male, 65-74 years				
...	All causes	218,165	100.0	2,516.2
1	Malignant neoplasms (C00-C97)	75,468	34.6	870.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	57,265	26.2	660.5
3	Chronic lower respiratory diseases (J40-J47)	14,152	6.5	163.2
4	Cerebrovascular diseases (I60-I69)	9,363	4.3	108.0
5	Diabetes mellitus (E10-E14)	8,344	3.8	96.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,212	2.4	60.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,909	1.8	45.1
8	Influenza and pneumonia (J10-J18)	3,371	1.5	38.9
9	Septicemia (A40-A41)	3,019	1.4	34.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	3,008	1.4	34.7
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,968	0.9	22.7
12	Alzheimer's disease (G30)	1,709	0.8	19.7
13	Aortic aneurysm and dissection (I71)	1,687	0.8	19.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,606	0.7	18.5
15	Parkinson's disease (G20-G21)	1,522	0.7	17.6
...	All other causes (Residual)	26,562	12.2	306.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 75-84 years				
...	All causes	327,315	100.0	6,177.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	92,378	28.2	1,743.5
2	Malignant neoplasms (C00-C97)	86,434	26.4	1,631.3
3	Chronic lower respiratory diseases (J40-J47)	23,167	7.1	437.3
4	Cerebrovascular diseases (I60-I69)	18,307	5.6	345.5
5	Diabetes mellitus (E10-E14)	10,286	3.1	194.1
6	Influenza and pneumonia (J10-J18)	8,315	2.5	156.9
7	Alzheimer's disease (G30)	8,295	2.5	156.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,246	2.2	136.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,194	2.2	135.8
10	Parkinson's disease (G20-G21)	5,546	1.7	104.7
11	Septicemia (A40-A41)	4,914	1.5	92.7
12	Pneumonitis due to solids and liquids (J69)	3,029	0.9	57.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,714	0.8	51.2
14	Aortic aneurysm and dissection (I71)	2,566	0.8	48.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,519	0.8	47.5
...	All other causes (Residual)	44,405	13.6	838.1
All races, male, 85 years and over				
...	All causes	241,578	100.0	14,309.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	81,373	33.7	4,819.9
2	Malignant neoplasms (C00-C97)	37,964	15.7	2,248.7
3	Cerebrovascular diseases (I60-I69)	15,742	6.5	932.4
4	Chronic lower respiratory diseases (J40-J47)	12,845	5.3	760.8
5	Alzheimer's disease (G30)	10,782	4.5	638.6
6	Influenza and pneumonia (J10-J18)	9,900	4.1	586.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,366	2.6	377.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,692	2.4	337.1
9	Diabetes mellitus (E10-E14)	5,419	2.2	321.0
10	Parkinson's disease (G20-G21)	3,911	1.6	231.7
11	Pneumonitis due to solids and liquids (J69)	3,796	1.6	224.8
12	Septicemia (A40-A41)	3,288	1.4	194.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,688	1.1	159.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,558	0.6	92.3
15	Aortic aneurysm and dissection (I71)	1,479	0.6	87.6
...	All other causes (Residual)	38,775	16.1	2,296.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, all ages				
...	All causes	1,224,322	100.0	806.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	315,930	25.8	208.0
2	Malignant neoplasms (C00-C97)	269,819	22.0	177.6
3	Cerebrovascular diseases (I60-I69)	82,595	6.7	54.4
4	Chronic lower respiratory diseases (J40-J47)	65,323	5.3	43.0
5	Alzheimer's disease (G30)	51,281	4.2	33.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42,658	3.5	28.1
7	Diabetes mellitus (E10-E14)	36,443	3.0	24.0
8	Influenza and pneumonia (J10-J18)	30,676	2.5	20.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23,250	1.9	15.3
10	Septicemia (A40-A41)	18,712	1.5	12.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14,440	1.2	9.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	9,689	0.8	6.4
13	Parkinson's disease (G20-G21)	8,266	0.7	5.4
14	Pneumonitis due to solids and liquids (J69)	7,971	0.7	5.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7,138	0.6	4.7
...	All other causes (Residual)	240,131	19.6	158.1
All races, female, 1-4 years				
...	All causes	2,090	100.0	26.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	683	32.7	8.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	271	13.0	3.4
3	Malignant neoplasms (C00-C97)	170	8.1	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	158	7.6	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	72	3.4	0.9
6	Influenza and pneumonia (J10-J18)	62	3.0	0.8
7	Septicemia (A40-A41)	45	2.2	0.6
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	1.7	0.5
9	Cerebrovascular diseases (I60-I69)	28	1.3	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	28	1.3	0.4
11	Meningitis (G00,G03)	14	0.7	*
11	Chronic lower respiratory diseases (J40-J47)	14	0.7	*
13	Anemias (D50-D64)	11	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	8	0.4	*
...	All other causes (Residual)	479	22.9	6.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 5-14 years				
...	All causes	2,524	100.0	12.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	873	34.6	4.4
2	Malignant neoplasms (C00-C97)	400	15.8	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	163	6.5	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	144	5.7	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	113	4.5	0.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	66	2.6	0.3
7	Cerebrovascular diseases (I60-I69)	39	1.5	0.2
8	Chronic lower respiratory diseases (J40-J47)	37	1.5	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	1.4	0.2
10	Influenza and pneumonia (J10-J18)	33	1.3	0.2
11	Septicemia (A40-A41)	32	1.3	0.2
12	Anemias (D50-D64)	18	0.7	*
13	Diabetes mellitus (E10-E14)	15	0.6	*
14	Certain conditions originating in the perinatal period (P00-P96)	10	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.3	*
...	All other causes (Residual)	538	21.3	2.7
All races, female, 15-24 years				
...	All causes	8,817	100.0	42.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,995	45.3	19.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	726	8.2	3.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	661	7.5	3.2
4	Malignant neoplasms (C00-C97)	637	7.2	3.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	365	4.1	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	181	2.1	0.9
7	Pregnancy, childbirth and the puerperium (O00-099)	179	2.0	0.9
8	Cerebrovascular diseases (I60-I69)	95	1.1	0.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	85	1.0	0.4
10	Influenza and pneumonia (J10-J18)	83	0.9	0.4
11	Septicemia (A40-A41)	71	0.8	0.3
12	Diabetes mellitus (E10-E14)	69	0.8	0.3
13	Chronic lower respiratory diseases (J40-J47)	58	0.7	0.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	49	0.6	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	0.5	0.2
...	All other causes (Residual)	1,520	17.2	7.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 25-34 years				
...	All causes	12,764	100.0	64.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,488	27.3	17.6
2	Malignant neoplasms (C00-C97)	1,886	14.8	9.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,009	7.9	5.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	932	7.3	4.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	742	5.8	3.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	453	3.5	2.3
7	Pregnancy, childbirth and the puerperium (O00-099)	324	2.5	1.6
8	Diabetes mellitus (E10-E14)	297	2.3	1.5
9	Cerebrovascular diseases (I60-I69)	237	1.9	1.2
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	174	1.4	0.9
11	Influenza and pneumonia (J10-J18)	160	1.3	0.8
12	Septicemia (A40-A41)	139	1.1	0.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	119	0.9	0.6
14	Chronic lower respiratory diseases (J40-J47)	118	0.9	0.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	101	0.8	0.5
...	All other causes (Residual)	2,585	20.3	13.0
All races, female, 35-44 years				
...	All causes	30,893	100.0	141.6
1	Malignant neoplasms (C00-C97)	7,938	25.7	36.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,103	16.5	23.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,714	12.0	17.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,526	4.9	7.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,181	3.8	5.4
6	Cerebrovascular diseases (I60-I69)	1,058	3.4	4.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	885	2.9	4.1
8	Diabetes mellitus (E10-E14)	815	2.6	3.7
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	751	2.4	3.4
10	Chronic lower respiratory diseases (J40-J47)	445	1.4	2.0
11	Septicemia (A40-A41)	418	1.4	1.9
12	Influenza and pneumonia (J10-J18)	379	1.2	1.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	327	1.1	1.5
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	203	0.7	0.9
15	Pregnancy, childbirth and the puerperium (O00-099)	192	0.6	0.9
...	All other causes (Residual)	5,958	19.3	27.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 45-54 years				
...	All causes	69,859	100.0	317.7
1	Malignant neoplasms (C00-C97)	24,993	35.8	113.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,662	15.3	48.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,838	8.4	26.6
4	Cerebrovascular diseases (I60-I69)	2,859	4.1	13.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,195	3.1	10.0
6	Diabetes mellitus (E10-E14)	2,177	3.1	9.9
7	Chronic lower respiratory diseases (J40-J47)	1,966	2.8	8.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,850	2.6	8.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,114	1.6	5.1
10	Septicemia (A40-A41)	1,013	1.5	4.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	907	1.3	4.1
12	Influenza and pneumonia (J10-J18)	821	1.2	3.7
13	Viral hepatitis (B15-B19)	809	1.2	3.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	581	0.8	2.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	486	0.7	2.2
...	All other causes (Residual)	11,588	16.6	52.7
All races, female, 55-64 years				
...	All causes	112,414	100.0	687.0
1	Malignant neoplasms (C00-C97)	46,106	41.0	281.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,300	18.1	124.1
3	Chronic lower respiratory diseases (J40-J47)	5,993	5.3	36.6
4	Diabetes mellitus (E10-E14)	4,762	4.2	29.1
5	Cerebrovascular diseases (I60-I69)	4,621	4.1	28.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,598	3.2	22.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,122	1.9	13.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,926	1.7	11.8
9	Septicemia (A40-A41)	1,888	1.7	11.5
10	Influenza and pneumonia (J10-J18)	1,277	1.1	7.8
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,120	1.0	6.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	905	0.8	5.5
13	Viral hepatitis (B15-B19)	625	0.6	3.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	583	0.5	3.6
15	Human immunodeficiency virus (HIV) disease (B20-B24)	370	0.3	2.3
...	All other causes (Residual)	16,218	14.4	99.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 65-74 years				
...	All causes	171,928	100.0	1,677.9
1	Malignant neoplasms (C00-C97)	62,086	36.1	605.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,487	20.6	346.3
3	Chronic lower respiratory diseases (J40-J47)	14,084	8.2	137.5
4	Cerebrovascular diseases (I60-I69)	8,860	5.2	86.5
5	Diabetes mellitus (E10-E14)	7,139	4.2	69.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,544	2.1	34.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,208	1.9	31.3
8	Septicemia (A40-A41)	3,045	1.8	29.7
9	Influenza and pneumonia (J10-J18)	2,690	1.6	26.3
10	Alzheimer's disease (G30)	2,103	1.2	20.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,909	1.1	18.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,568	0.9	15.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,009	0.6	9.8
14	Aortic aneurysm and dissection (I71)	948	0.6	9.3
15	Parkinson's disease (G20-G21)	788	0.5	7.7
...	All other causes (Residual)	23,460	13.6	229.0
All races, female, 75-84 years				
...	All causes	340,023	100.0	4,388.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	88,073	25.9	1,136.7
2	Malignant neoplasms (C00-C97)	78,455	23.1	1,012.5
3	Cerebrovascular diseases (I60-I69)	25,412	7.5	328.0
4	Chronic lower respiratory diseases (J40-J47)	24,239	7.1	312.8
5	Alzheimer's disease (G30)	14,620	4.3	188.7
6	Diabetes mellitus (E10-E14)	11,477	3.4	148.1
7	Influenza and pneumonia (J10-J18)	8,353	2.5	107.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,290	2.1	94.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,514	1.9	84.1
10	Septicemia (A40-A41)	5,831	1.7	75.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,940	1.2	50.8
12	Parkinson's disease (G20-G21)	3,555	1.0	45.9
13	Pneumonitis due to solids and liquids (J69)	2,255	0.7	29.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,231	0.7	28.8
15	Aortic aneurysm and dissection (I71)	2,056	0.6	26.5
...	All other causes (Residual)	55,722	16.4	719.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 85 years and over				
...	All causes	460,414	100.0	12,759.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	155,966	33.9	4,322.1
2	Malignant neoplasms (C00-C97)	47,108	10.2	1,305.5
3	Cerebrovascular diseases (I60-I69)	39,326	8.5	1,089.8
4	Alzheimer's disease (G30)	34,151	7.4	946.4
5	Chronic lower respiratory diseases (J40-J47)	18,358	4.0	508.7
6	Influenza and pneumonia (J10-J18)	16,717	3.6	463.3
7	Diabetes mellitus (E10-E14)	9,686	2.1	268.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9,022	2.0	250.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,869	1.9	245.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,342	1.6	203.5
11	Septicemia (A40-A41)	6,104	1.3	169.2
12	Pneumonitis due to solids and liquids (J69)	4,458	1.0	123.5
13	Parkinson's disease (G20-G21)	3,758	0.8	104.1
14	Atherosclerosis (I70)	3,324	0.7	92.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,576	0.6	71.4
...	All other causes (Residual)	93,649	20.3	2,595.2
White, both sexes, all ages				
...	All causes	2,077,549	100.0	858.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	545,974	26.3	225.5
2	Malignant neoplasms (C00-C97)	482,575	23.2	199.3
3	Cerebrovascular diseases (I60-I69)	115,864	5.6	47.9
4	Chronic lower respiratory diseases (J40-J47)	114,993	5.5	47.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	103,853	5.0	42.9
6	Alzheimer's disease (G30)	67,088	3.2	27.7
7	Diabetes mellitus (E10-E14)	57,204	2.8	23.6
8	Influenza and pneumonia (J10-J18)	49,401	2.4	20.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35,793	1.7	14.8
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30,138	1.5	12.4
11	Septicemia (A40-A41)	27,373	1.3	11.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	24,191	1.2	10.0
13	Parkinson's disease (G20-G21)	18,517	0.9	7.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18,264	0.9	7.5
15	Pneumonitis due to solids and liquids (J69)	15,136	0.7	6.3
...	All other causes (Residual)	371,185	17.9	153.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 1-4 years				
...	All causes	3,233	100.0	25.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,170	36.2	9.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	376	11.6	3.0
3	Malignant neoplasms (C00-C97)	286	8.8	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	190	5.9	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	110	3.4	0.9
6	Influenza and pneumonia (J10-J18)	83	2.6	0.7
7	Septicemia (A40-A41)	59	1.8	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	1.6	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	40	1.2	0.3
10	Cerebrovascular diseases (I60-I69)	39	1.2	0.3
11	Chronic lower respiratory diseases (J40-J47)	26	0.8	0.2
12	Complications of medical and surgical care (Y40-Y84,Y88)	17	0.5	*
13	Meningitis (G00,G03)	15	0.5	*
14	Meningococcal infection (A39)	13	0.4	*
14	Anemias (D50-D64)	13	0.4	*
...	All other causes (Residual)	745	23.0	5.9
White, both sexes, 5-14 years				
...	All causes	4,460	100.0	14.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,668	37.4	5.3
2	Malignant neoplasms (C00-C97)	720	16.1	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	259	5.8	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	227	5.1	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	170	3.8	0.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	167	3.7	0.5
7	Cerebrovascular diseases (I60-I69)	67	1.5	0.2
8	Septicemia (A40-A41)	63	1.4	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	59	1.3	0.2
10	Influenza and pneumonia (J10-J18)	53	1.2	0.2
11	Chronic lower respiratory diseases (J40-J47)	42	0.9	0.1
12	Diabetes mellitus (E10-E14)	19	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	18	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	14	0.3	*
15	Anemias (D50-D64)	13	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.3	*
...	All other causes (Residual)	888	19.9	2.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 15-24 years				
...	All causes	25,871	100.0	78.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,824	53.4	41.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,477	13.4	10.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,213	8.6	6.7
4	Malignant neoplasms (C00-C97)	1,294	5.0	3.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	730	2.8	2.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	343	1.3	1.0
7	Cerebrovascular diseases (I60-I69)	159	0.6	0.5
8	Influenza and pneumonia (J10-J18)	138	0.5	0.4
9	Pregnancy, childbirth and the puerperium (O00-099)	116	0.4	0.3
10	Diabetes mellitus (E10-E14)	106	0.4	0.3
11	Septicemia (A40-A41)	90	0.3	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	70	0.3	0.2
13	Chronic lower respiratory diseases (J40-J47)	69	0.3	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	66	0.3	0.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	59	0.2	0.2
...	All other causes (Residual)	3,117	12.0	9.4
White, both sexes, 25-34 years				
...	All causes	31,127	100.0	98.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,473	40.1	39.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,259	13.7	13.4
3	Malignant neoplasms (C00-C97)	2,799	9.0	8.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,143	6.9	6.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,902	6.1	6.0
6	Diabetes mellitus (E10-E14)	406	1.3	1.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	405	1.3	1.3
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	349	1.1	1.1
9	Cerebrovascular diseases (I60-I69)	345	1.1	1.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	250	0.8	0.8
11	Influenza and pneumonia (J10-J18)	238	0.8	0.8
12	Septicemia (A40-A41)	186	0.6	0.6
13	Pregnancy, childbirth and the puerperium (O00-099)	180	0.6	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	158	0.5	0.5
15	Chronic lower respiratory diseases (J40-J47)	141	0.5	0.4
...	All other causes (Residual)	4,893	15.7	15.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 35-44 years				
...	All causes	62,060	100.0	177.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,647	23.6	41.8
2	Malignant neoplasms (C00-C97)	10,755	17.3	30.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,879	14.3	25.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,945	9.6	17.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,136	3.4	6.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,760	2.8	5.0
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,569	2.5	4.5
8	Diabetes mellitus (E10-E14)	1,441	2.3	4.1
9	Cerebrovascular diseases (I60-I69)	1,389	2.2	4.0
10	Influenza and pneumonia (J10-J18)	628	1.0	1.8
11	Chronic lower respiratory diseases (J40-J47)	591	1.0	1.7
12	Septicemia (A40-A41)	572	0.9	1.6
13	Viral hepatitis (B15-B19)	421	0.7	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	411	0.7	1.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	381	0.6	1.1
...	All other causes (Residual)	10,535	17.0	30.1
White, both sexes, 45-54 years				
...	All causes	141,969	100.0	397.9
1	Malignant neoplasms (C00-C97)	39,795	28.0	111.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,462	20.0	79.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,245	11.4	45.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,924	4.9	19.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	6,610	4.7	18.5
6	Cerebrovascular diseases (I60-I69)	4,135	2.9	11.6
7	Diabetes mellitus (E10-E14)	4,059	2.9	11.4
8	Chronic lower respiratory diseases (J40-J47)	3,236	2.3	9.1
9	Viral hepatitis (B15-B19)	2,415	1.7	6.8
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,874	1.3	5.3
11	Septicemia (A40-A41)	1,509	1.1	4.2
12	Influenza and pneumonia (J10-J18)	1,492	1.1	4.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,334	0.9	3.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,283	0.9	3.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	693	0.5	1.9
...	All other causes (Residual)	21,903	15.4	61.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 55-64 years				
...	All causes	227,108	100.0	847.5
1	Malignant neoplasms (C00-C97)	84,421	37.2	315.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	51,715	22.8	193.0
3	Chronic lower respiratory diseases (J40-J47)	11,000	4.8	41.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,514	4.2	35.5
5	Diabetes mellitus (E10-E14)	8,482	3.7	31.7
6	Cerebrovascular diseases (I60-I69)	7,417	3.3	27.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	6,285	2.8	23.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,322	1.9	16.1
9	Septicemia (A40-A41)	2,929	1.3	10.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,875	1.3	10.7
11	Influenza and pneumonia (J10-J18)	2,557	1.1	9.5
12	Viral hepatitis (B15-B19)	1,626	0.7	6.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,361	0.6	5.1
14	Aortic aneurysm and dissection (I71)	1,122	0.5	4.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,109	0.5	4.1
...	All other causes (Residual)	30,373	13.4	113.3
White, both sexes, 65-74 years				
...	All causes	328,578	100.0	2,021.1
1	Malignant neoplasms (C00-C97)	118,280	36.0	727.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	76,373	23.2	469.8
3	Chronic lower respiratory diseases (J40-J47)	26,055	7.9	160.3
4	Cerebrovascular diseases (I60-I69)	14,230	4.3	87.5
5	Diabetes mellitus (E10-E14)	11,915	3.6	73.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,266	2.2	44.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,357	1.6	33.0
8	Influenza and pneumonia (J10-J18)	5,061	1.5	31.1
9	Septicemia (A40-A41)	4,668	1.4	28.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	4,424	1.3	27.2
11	Alzheimer's disease (G30)	3,433	1.0	21.1
12	Aortic aneurysm and dissection (I71)	2,302	0.7	14.2
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,243	0.7	13.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,180	0.7	13.4
15	Parkinson's disease (G20-G21)	2,120	0.6	13.0
...	All other causes (Residual)	42,671	13.0	262.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 75-84 years				
...	All causes	593,016	100.0	5,127.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	159,346	26.9	1,377.8
2	Malignant neoplasms (C00-C97)	147,150	24.8	1,272.3
3	Chronic lower respiratory diseases (J40-J47)	44,390	7.5	383.8
4	Cerebrovascular diseases (I60-I69)	37,984	6.4	328.4
5	Alzheimer's disease (G30)	21,126	3.6	182.7
6	Diabetes mellitus (E10-E14)	17,898	3.0	154.8
7	Influenza and pneumonia (J10-J18)	14,750	2.5	127.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,529	2.1	108.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12,060	2.0	104.3
10	Septicemia (A40-A41)	9,001	1.5	77.8
11	Parkinson's disease (G20-G21)	8,641	1.5	74.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,266	0.9	45.5
13	Pneumonitis due to solids and liquids (J69)	4,751	0.8	41.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,335	0.7	37.5
15	Aortic aneurysm and dissection (I71)	4,213	0.7	36.4
...	All other causes (Residual)	89,576	15.1	774.5
White, both sexes, 85 years and over				
...	All causes	641,545	100.0	13,480.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	217,804	33.9	4,576.6
2	Malignant neoplasms (C00-C97)	77,005	12.0	1,618.1
3	Cerebrovascular diseases (I60-I69)	50,009	7.8	1,050.8
4	Alzheimer's disease (G30)	41,827	6.5	878.9
5	Chronic lower respiratory diseases (J40-J47)	29,425	4.6	618.3
6	Influenza and pneumonia (J10-J18)	24,237	3.8	509.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,737	2.1	288.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13,395	2.1	281.5
9	Diabetes mellitus (E10-E14)	12,867	2.0	270.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8,542	1.3	179.5
11	Septicemia (A40-A41)	8,140	1.3	171.0
12	Pneumonitis due to solids and liquids (J69)	7,616	1.2	160.0
13	Parkinson's disease (G20-G21)	7,332	1.1	154.1
14	Atherosclerosis (I70)	4,435	0.7	93.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3,809	0.6	80.0
...	All other causes (Residual)	121,365	18.9	2,550.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, all ages				
...	All causes	1,022,328	100.0	852.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	272,117	26.6	226.9
2	Malignant neoplasms (C00-C97)	250,322	24.5	208.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	66,843	6.5	55.7
4	Chronic lower respiratory diseases (J40-J47)	54,043	5.3	45.1
5	Cerebrovascular diseases (I60-I69)	45,198	4.4	37.7
6	Diabetes mellitus (E10-E14)	29,060	2.8	24.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23,767	2.3	19.8
8	Influenza and pneumonia (J10-J18)	22,310	2.2	18.6
9	Alzheimer's disease (G30)	19,654	1.9	16.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,715	1.7	14.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	15,739	1.5	13.1
12	Septicemia (A40-A41)	12,450	1.2	10.4
13	Parkinson's disease (G20-G21)	10,706	1.0	8.9
14	Pneumonitis due to solids and liquids (J69)	7,984	0.8	6.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6,990	0.7	5.8
...	All other causes (Residual)	167,430	16.4	139.6
White, male, 1-4 years				
...	All causes	1,781	100.0	27.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	675	37.9	10.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	191	10.7	2.9
3	Malignant neoplasms (C00-C97)	162	9.1	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	109	6.1	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	60	3.4	0.9
6	Influenza and pneumonia (J10-J18)	43	2.4	0.7
7	Septicemia (A40-A41)	29	1.6	0.4
8	Certain conditions originating in the perinatal period (P00-P96)	26	1.5	0.4
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.1	*
10	Chronic lower respiratory diseases (J40-J47)	17	1.0	*
11	Cerebrovascular diseases (I60-I69)	16	0.9	*
12	Meningococcal infection (A39)	9	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.4	*
15	Anemias (D50-D64)	5	0.3	*
...	All other causes (Residual)	404	22.7	6.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 5-14 years				
...	All causes	2,639	100.0	16.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,017	38.5	6.3
2	Malignant neoplasms (C00-C97)	401	15.2	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	148	5.6	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	130	4.9	0.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	117	4.4	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	99	3.8	0.6
7	Cerebrovascular diseases (I60-I69)	46	1.7	0.3
8	Septicemia (A40-A41)	38	1.4	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	1.3	0.2
10	Chronic lower respiratory diseases (J40-J47)	32	1.2	0.2
11	Influenza and pneumonia (J10-J18)	31	1.2	0.2
12	Certain conditions originating in the perinatal period (P00-P96)	12	0.5	*
13	Diabetes mellitus (E10-E14)	11	0.4	*
14	Pneumonitis due to solids and liquids (J69)	8	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.3	*
...	All other causes (Residual)	500	18.9	3.1
White, male, 15-24 years				
...	All causes	19,175	100.0	111.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,423	54.4	60.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,939	15.3	17.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,824	9.5	10.6
4	Malignant neoplasms (C00-C97)	805	4.2	4.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	505	2.6	2.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	206	1.1	1.2
7	Cerebrovascular diseases (I60-I69)	91	0.5	0.5
8	Influenza and pneumonia (J10-J18)	75	0.4	0.4
9	Diabetes mellitus (E10-E14)	60	0.3	0.3
10	Legal intervention (Y35,Y89.0)	50	0.3	0.3
11	Septicemia (A40-A41)	43	0.2	0.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	0.2	0.3
13	Chronic lower respiratory diseases (J40-J47)	40	0.2	0.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	0.2	0.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	36	0.2	0.2
...	All other causes (Residual)	1,996	10.4	11.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 25-34 years				
...	All causes	22,083	100.0	135.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,564	43.3	58.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,441	15.6	21.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,537	7.0	9.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,520	6.9	9.3
5	Malignant neoplasms (C00-C97)	1,396	6.3	8.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	305	1.4	1.9
7	Diabetes mellitus (E10-E14)	224	1.0	1.4
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	212	1.0	1.3
9	Cerebrovascular diseases (I60-I69)	195	0.9	1.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	174	0.8	1.1
11	Influenza and pneumonia (J10-J18)	128	0.6	0.8
12	Septicemia (A40-A41)	99	0.4	0.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	95	0.4	0.6
14	Legal intervention (Y35,Y89.0)	89	0.4	0.5
15	Chronic lower respiratory diseases (J40-J47)	70	0.3	0.4
...	All other causes (Residual)	3,034	13.7	18.6
White, male, 35-44 years				
...	All causes	39,770	100.0	224.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,393	26.1	58.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,437	16.2	36.3
3	Malignant neoplasms (C00-C97)	4,726	11.9	26.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,548	11.4	25.7
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,427	3.6	8.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,424	3.6	8.0
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,131	2.8	6.4
8	Diabetes mellitus (E10-E14)	894	2.2	5.0
9	Cerebrovascular diseases (I60-I69)	749	1.9	4.2
10	Influenza and pneumonia (J10-J18)	349	0.9	2.0
11	Septicemia (A40-A41)	306	0.8	1.7
12	Viral hepatitis (B15-B19)	291	0.7	1.6
13	Chronic lower respiratory diseases (J40-J47)	282	0.7	1.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	252	0.6	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	218	0.5	1.2
...	All other causes (Residual)	6,343	15.9	35.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 45-54 years				
...	All causes	89,681	100.0	505.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,158	23.6	119.2
2	Malignant neoplasms (C00-C97)	20,163	22.5	113.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,380	12.7	64.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,192	5.8	29.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	4,772	5.3	26.9
6	Diabetes mellitus (E10-E14)	2,583	2.9	14.6
7	Cerebrovascular diseases (I60-I69)	2,279	2.5	12.8
8	Viral hepatitis (B15-B19)	1,770	2.0	10.0
9	Chronic lower respiratory diseases (J40-J47)	1,604	1.8	9.0
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,573	1.8	8.9
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	910	1.0	5.1
12	Influenza and pneumonia (J10-J18)	874	1.0	4.9
13	Septicemia (A40-A41)	856	1.0	4.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	820	0.9	4.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	458	0.5	2.6
...	All other causes (Residual)	13,289	14.8	74.9
White, male, 55-64 years				
...	All causes	137,161	100.0	1,050.6
1	Malignant neoplasms (C00-C97)	46,069	33.6	352.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,407	26.5	278.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,483	4.7	49.7
4	Chronic lower respiratory diseases (J40-J47)	5,595	4.1	42.9
5	Diabetes mellitus (E10-E14)	5,094	3.7	39.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,464	3.3	34.2
7	Cerebrovascular diseases (I60-I69)	4,110	3.0	31.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,259	2.4	25.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,659	1.2	12.7
10	Septicemia (A40-A41)	1,549	1.1	11.9
11	Influenza and pneumonia (J10-J18)	1,522	1.1	11.7
12	Viral hepatitis (B15-B19)	1,154	0.8	8.8
13	Aortic aneurysm and dissection (I71)	843	0.6	6.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	827	0.6	6.3
15	Human immunodeficiency virus (HIV) disease (B20-B24)	631	0.5	4.8
...	All other causes (Residual)	17,495	12.8	134.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 65-74 years				
...	All causes	184,925	100.0	2,455.8
1	Malignant neoplasms (C00-C97)	64,913	35.1	862.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	47,936	25.9	636.6
3	Chronic lower respiratory diseases (J40-J47)	12,916	7.0	171.5
4	Cerebrovascular diseases (I60-I69)	7,312	4.0	97.1
5	Diabetes mellitus (E10-E14)	6,658	3.6	88.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,502	2.4	59.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,918	1.6	38.8
8	Influenza and pneumonia (J10-J18)	2,810	1.5	37.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	2,708	1.5	36.0
10	Septicemia (A40-A41)	2,373	1.3	31.5
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,862	1.0	24.7
12	Alzheimer's disease (G30)	1,546	0.8	20.5
13	Aortic aneurysm and dissection (I71)	1,499	0.8	19.9
14	Parkinson's disease (G20-G21)	1,398	0.8	18.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,215	0.7	16.1
...	All other causes (Residual)	22,359	12.1	296.9
White, male, 75-84 years				
...	All causes	293,004	100.0	6,182.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	82,625	28.2	1,743.3
2	Malignant neoplasms (C00-C97)	77,316	26.4	1,631.3
3	Chronic lower respiratory diseases (J40-J47)	21,486	7.3	453.3
4	Cerebrovascular diseases (I60-I69)	16,041	5.5	338.5
5	Diabetes mellitus (E10-E14)	8,785	3.0	185.4
6	Alzheimer's disease (G30)	7,701	2.6	162.5
7	Influenza and pneumonia (J10-J18)	7,382	2.5	155.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,584	2.2	138.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,157	2.1	129.9
10	Parkinson's disease (G20-G21)	5,263	1.8	111.0
11	Septicemia (A40-A41)	4,174	1.4	88.1
12	Pneumonitis due to solids and liquids (J69)	2,726	0.9	57.5
13	Aortic aneurysm and dissection (I71)	2,386	0.8	50.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,322	0.8	49.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,164	0.7	45.7
...	All other causes (Residual)	39,892	13.6	841.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 85 years and over				
...	All causes	221,622	100.0	14,576.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	75,215	33.9	4,947.1
2	Malignant neoplasms (C00-C97)	34,334	15.5	2,258.3
3	Cerebrovascular diseases (I60-I69)	14,311	6.5	941.3
4	Chronic lower respiratory diseases (J40-J47)	11,989	5.4	788.6
5	Alzheimer's disease (G30)	10,072	4.5	662.5
6	Influenza and pneumonia (J10-J18)	8,991	4.1	591.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,683	2.6	373.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,358	2.4	352.4
9	Diabetes mellitus (E10-E14)	4,743	2.1	312.0
10	Parkinson's disease (G20-G21)	3,760	1.7	247.3
11	Pneumonitis due to solids and liquids (J69)	3,524	1.6	231.8
12	Septicemia (A40-A41)	2,899	1.3	190.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,300	1.0	151.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,425	0.6	93.7
15	Aortic aneurysm and dissection (I71)	1,375	0.6	90.4
...	All other causes (Residual)	35,643	16.1	2,344.4
White, female, all ages				
...	All causes	1,055,221	100.0	863.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	273,857	26.0	224.2
2	Malignant neoplasms (C00-C97)	232,253	22.0	190.1
3	Cerebrovascular diseases (I60-I69)	70,666	6.7	57.9
4	Chronic lower respiratory diseases (J40-J47)	60,950	5.8	49.9
5	Alzheimer's disease (G30)	47,434	4.5	38.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37,010	3.5	30.3
7	Diabetes mellitus (E10-E14)	28,144	2.7	23.0
8	Influenza and pneumonia (J10-J18)	27,091	2.6	22.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,078	1.7	14.8
10	Septicemia (A40-A41)	14,923	1.4	12.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,274	1.1	9.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	8,452	0.8	6.9
13	Parkinson's disease (G20-G21)	7,811	0.7	6.4
14	Pneumonitis due to solids and liquids (J69)	7,152	0.7	5.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,371	0.6	5.2
...	All other causes (Residual)	203,755	19.3	166.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 1-4 years				
...	All causes	1,452	100.0	23.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	495	34.1	8.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	185	12.7	3.0
3	Malignant neoplasms (C00-C97)	124	8.5	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	5.6	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	50	3.4	0.8
6	Influenza and pneumonia (J10-J18)	40	2.8	0.6
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	2.2	0.5
8	Septicemia (A40-A41)	30	2.1	0.5
9	Cerebrovascular diseases (I60-I69)	23	1.6	0.4
10	Certain conditions originating in the perinatal period (P00-P96)	14	1.0	*
11	Meningitis (G00,G03)	11	0.8	*
12	Chronic lower respiratory diseases (J40-J47)	9	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.6	*
14	Anemias (D50-D64)	8	0.6	*
15	Acute bronchitis and bronchiolitis (J20-J21)	5	0.3	*
15	Pneumonitis due to solids and liquids (J69)	5	0.3	*
...	All other causes (Residual)	331	22.8	5.4
White, female, 5-14 years				
...	All causes	1,821	100.0	11.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	651	35.7	4.3
2	Malignant neoplasms (C00-C97)	319	17.5	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	111	6.1	0.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	97	5.3	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	71	3.9	0.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	50	2.7	0.3
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	1.4	0.2
8	Septicemia (A40-A41)	25	1.4	0.2
9	Influenza and pneumonia (J10-J18)	22	1.2	0.1
10	Cerebrovascular diseases (I60-I69)	21	1.2	0.1
11	Chronic lower respiratory diseases (J40-J47)	10	0.5	*
12	Diabetes mellitus (E10-E14)	8	0.4	*
13	Anemias (D50-D64)	7	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	6	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.3	*
...	All other causes (Residual)	391	21.5	2.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 15-24 years				
...	All causes	6,696	100.0	41.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,401	50.8	21.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	538	8.0	3.4
3	Malignant neoplasms (C00-C97)	489	7.3	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	389	5.8	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	225	3.4	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	137	2.0	0.9
7	Pregnancy, childbirth and the puerperium (O00-099)	116	1.7	0.7
8	Cerebrovascular diseases (I60-I69)	68	1.0	0.4
9	Influenza and pneumonia (J10-J18)	63	0.9	0.4
10	Septicemia (A40-A41)	47	0.7	0.3
11	Diabetes mellitus (E10-E14)	46	0.7	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	0.5	0.2
13	Chronic lower respiratory diseases (J40-J47)	29	0.4	0.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	23	0.3	0.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.3	0.1
...	All other causes (Residual)	1,071	16.0	6.7
White, female, 25-34 years				
...	All causes	9,044	100.0	58.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,909	32.2	18.9
2	Malignant neoplasms (C00-C97)	1,403	15.5	9.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	818	9.0	5.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	606	6.7	3.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	382	4.2	2.5
6	Diabetes mellitus (E10-E14)	182	2.0	1.2
7	Pregnancy, childbirth and the puerperium (O00-099)	180	2.0	1.2
8	Cerebrovascular diseases (I60-I69)	150	1.7	1.0
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	137	1.5	0.9
10	Influenza and pneumonia (J10-J18)	110	1.2	0.7
11	Human immunodeficiency virus (HIV) disease (B20-B24)	100	1.1	0.7
12	Septicemia (A40-A41)	87	1.0	0.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	76	0.8	0.5
14	Chronic lower respiratory diseases (J40-J47)	71	0.8	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	63	0.7	0.4
...	All other causes (Residual)	1,770	19.6	11.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 35-44 years				
...	All causes	22,290	100.0	129.0
1	Malignant neoplasms (C00-C97)	6,029	27.0	34.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,254	19.1	24.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,442	11.0	14.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,397	6.3	8.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	712	3.2	4.1
6	Cerebrovascular diseases (I60-I69)	640	2.9	3.7
7	Diabetes mellitus (E10-E14)	547	2.5	3.2
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	438	2.0	2.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	333	1.5	1.9
10	Chronic lower respiratory diseases (J40-J47)	309	1.4	1.8
11	Influenza and pneumonia (J10-J18)	279	1.3	1.6
12	Septicemia (A40-A41)	266	1.2	1.5
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	163	0.7	0.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	159	0.7	0.9
15	Viral hepatitis (B15-B19)	130	0.6	0.8
...	All other causes (Residual)	4,192	18.8	24.3
White, female, 45-54 years				
...	All causes	52,288	100.0	291.6
1	Malignant neoplasms (C00-C97)	19,632	37.5	109.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,304	14.0	40.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,865	9.3	27.1
4	Cerebrovascular diseases (I60-I69)	1,856	3.5	10.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,838	3.5	10.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,732	3.3	9.7
7	Chronic lower respiratory diseases (J40-J47)	1,632	3.1	9.1
8	Diabetes mellitus (E10-E14)	1,476	2.8	8.2
9	Septicemia (A40-A41)	653	1.2	3.6
10	Viral hepatitis (B15-B19)	645	1.2	3.6
11	Influenza and pneumonia (J10-J18)	618	1.2	3.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	514	1.0	2.9
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	373	0.7	2.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	301	0.6	1.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	277	0.5	1.5
...	All other causes (Residual)	8,572	16.4	47.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 55-64 years				
...	All causes	89,947	100.0	654.6
1	Malignant neoplasms (C00-C97)	38,352	42.6	279.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,308	17.0	111.4
3	Chronic lower respiratory diseases (J40-J47)	5,405	6.0	39.3
4	Diabetes mellitus (E10-E14)	3,388	3.8	24.7
5	Cerebrovascular diseases (I60-I69)	3,307	3.7	24.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,031	3.4	22.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,821	2.0	13.3
8	Septicemia (A40-A41)	1,380	1.5	10.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,216	1.4	8.8
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,063	1.2	7.7
11	Influenza and pneumonia (J10-J18)	1,035	1.2	7.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	534	0.6	3.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	480	0.5	3.5
14	Viral hepatitis (B15-B19)	472	0.5	3.4
15	Alzheimer's disease (G30)	320	0.4	2.3
...	All other causes (Residual)	12,835	14.3	93.4
White, female, 65-74 years				
...	All causes	143,653	100.0	1,646.0
1	Malignant neoplasms (C00-C97)	53,367	37.1	611.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,437	19.8	325.8
3	Chronic lower respiratory diseases (J40-J47)	13,139	9.1	150.6
4	Cerebrovascular diseases (I60-I69)	6,918	4.8	79.3
5	Diabetes mellitus (E10-E14)	5,257	3.7	60.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,764	1.9	31.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,439	1.7	27.9
8	Septicemia (A40-A41)	2,295	1.6	26.3
9	Influenza and pneumonia (J10-J18)	2,251	1.6	25.8
10	Alzheimer's disease (G30)	1,887	1.3	21.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,716	1.2	19.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,082	0.8	12.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	874	0.6	10.0
14	Aortic aneurysm and dissection (I71)	803	0.6	9.2
15	Parkinson's disease (G20-G21)	722	0.5	8.3
...	All other causes (Residual)	19,702	13.7	225.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 75-84 years				
...	All causes	300,012	100.0	4,395.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	76,721	25.6	1,123.9
2	Malignant neoplasms (C00-C97)	69,834	23.3	1,023.0
3	Chronic lower respiratory diseases (J40-J47)	22,904	7.6	335.5
4	Cerebrovascular diseases (I60-I69)	21,943	7.3	321.5
5	Alzheimer's disease (G30)	13,425	4.5	196.7
6	Diabetes mellitus (E10-E14)	9,113	3.0	133.5
7	Influenza and pneumonia (J10-J18)	7,368	2.5	107.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,945	2.0	87.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,903	2.0	86.5
10	Septicemia (A40-A41)	4,827	1.6	70.7
11	Parkinson's disease (G20-G21)	3,378	1.1	49.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,102	1.0	45.4
13	Pneumonitis due to solids and liquids (J69)	2,025	0.7	29.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,013	0.7	29.5
15	Aortic aneurysm and dissection (I71)	1,827	0.6	26.8
...	All other causes (Residual)	49,684	16.6	727.9
White, female, 85 years and over				
...	All causes	419,923	100.0	12,965.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	142,589	34.0	4,402.6
2	Malignant neoplasms (C00-C97)	42,671	10.2	1,317.5
3	Cerebrovascular diseases (I60-I69)	35,698	8.5	1,102.2
4	Alzheimer's disease (G30)	31,755	7.6	980.5
5	Chronic lower respiratory diseases (J40-J47)	17,436	4.2	538.4
6	Influenza and pneumonia (J10-J18)	15,246	3.6	470.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,379	2.0	258.7
8	Diabetes mellitus (E10-E14)	8,124	1.9	250.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,712	1.8	238.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6,242	1.5	192.7
11	Septicemia (A40-A41)	5,241	1.2	161.8
12	Pneumonitis due to solids and liquids (J69)	4,092	1.0	126.3
13	Parkinson's disease (G20-G21)	3,572	0.9	110.3
14	Atherosclerosis (I70)	3,098	0.7	95.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,384	0.6	73.6
...	All other causes (Residual)	85,684	20.4	2,645.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, all ages				
...	All causes	289,971	100.0	733.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	72,253	24.9	182.6
2	Malignant neoplasms (C00-C97)	63,082	21.8	159.5
3	Cerebrovascular diseases (I60-I69)	17,045	5.9	43.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,917	4.8	35.2
5	Diabetes mellitus (E10-E14)	12,813	4.4	32.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9,032	3.1	22.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,397	2.9	21.2
8	Chronic lower respiratory diseases (J40-J47)	7,730	2.7	19.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	6,854	2.4	17.3
10	Septicemia (A40-A41)	6,108	2.1	15.4
11	Influenza and pneumonia (J10-J18)	5,311	1.8	13.4
12	Certain conditions originating in the perinatal period (P00-P96)	5,121	1.8	12.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,900	1.7	12.4
14	Alzheimer's disease (G30)	4,455	1.5	11.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,349	0.8	5.9
...	All other causes (Residual)	50,604	17.5	127.9
Black, both sexes, 1-4 years				
...	All causes	1,144	100.0	43.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	371	32.4	14.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	154	13.5	5.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	107	9.4	4.1
4	Malignant neoplasms (C00-C97)	65	5.7	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	44	3.8	1.7
6	Influenza and pneumonia (J10-J18)	34	3.0	1.3
7	Certain conditions originating in the perinatal period (P00-P96)	22	1.9	0.8
8	Septicemia (A40-A41)	21	1.8	0.8
9	Chronic lower respiratory diseases (J40-J47)	15	1.3	*
10	Anemias (D50-D64)	12	1.0	*
10	Cerebrovascular diseases (I60-I69)	12	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.8	*
13	Acute bronchitis and bronchiolitis (J20-J21)	5	0.4	*
14	Meningitis (G00,G03)	4	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.3	*
...	All other causes (Residual)	265	23.2	10.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 5-14 years				
...	All causes	1,394	100.0	21.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	471	33.8	7.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	149	10.7	2.3
3	Malignant neoplasms (C00-C97)	145	10.4	2.2
4	Chronic lower respiratory diseases (J40-J47)	69	4.9	1.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	67	4.8	1.0
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	67	4.8	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	2.7	0.6
8	Cerebrovascular diseases (I60-I69)	21	1.5	0.3
9	Septicemia (A40-A41)	20	1.4	0.3
9	Anemias (D50-D64)	20	1.4	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	1.1	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	11	0.8	*
13	Diabetes mellitus (E10-E14)	9	0.6	*
13	Influenza and pneumonia (J10-J18)	9	0.6	*
15	Meningitis (G00,G03)	8	0.6	*
...	All other causes (Residual)	276	19.8	4.2
Black, both sexes, 15-24 years				
...	All causes	7,483	100.0	112.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,295	44.0	49.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,750	23.4	26.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	416	5.6	6.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	308	4.1	4.6
5	Malignant neoplasms (C00-C97)	264	3.5	4.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	144	1.9	2.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	88	1.2	1.3
8	Chronic lower respiratory diseases (J40-J47)	75	1.0	1.1
9	Anemias (D50-D64)	64	0.9	1.0
10	Pregnancy, childbirth and the puerperium (O00-099)	58	0.8	0.9
11	Diabetes mellitus (E10-E14)	55	0.7	0.8
12	Cerebrovascular diseases (I60-I69)	44	0.6	0.7
13	Septicemia (A40-A41)	41	0.5	0.6
13	Influenza and pneumonia (J10-J18)	41	0.5	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	0.4	0.4
...	All other causes (Residual)	810	10.8	12.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 25-34 years				
...	All causes	10,045	100.0	177.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,670	26.6	47.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,933	19.2	34.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,033	10.3	18.3
4	Human immunodeficiency virus (HIV) disease (B20-B24)	770	7.7	13.6
5	Malignant neoplasms (C00-C97)	639	6.4	11.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	474	4.7	8.4
7	Diabetes mellitus (E10-E14)	240	2.4	4.2
8	Cerebrovascular diseases (I60-I69)	151	1.5	2.7
9	Pregnancy, childbirth and the puerperium (O00-099)	118	1.2	2.1
10	Anemias (D50-D64)	112	1.1	2.0
11	Chronic lower respiratory diseases (J40-J47)	108	1.1	1.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	107	1.1	1.9
13	Septicemia (A40-A41)	89	0.9	1.6
14	Influenza and pneumonia (J10-J18)	77	0.8	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	73	0.7	1.3
...	All other causes (Residual)	1,451	14.4	25.7
Black, both sexes, 35-44 years				
...	All causes	17,930	100.0	314.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,078	17.2	53.9
2	Malignant neoplasms (C00-C97)	2,515	14.0	44.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,347	13.1	41.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	2,189	12.2	38.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,347	7.5	23.6
6	Cerebrovascular diseases (I60-I69)	697	3.9	12.2
7	Diabetes mellitus (E10-E14)	583	3.3	10.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	418	2.3	7.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	336	1.9	5.9
10	Septicemia (A40-A41)	273	1.5	4.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	256	1.4	4.5
12	Chronic lower respiratory diseases (J40-J47)	226	1.3	4.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	209	1.2	3.7
14	Influenza and pneumonia (J10-J18)	186	1.0	3.3
15	Anemias (D50-D64)	146	0.8	2.6
...	All other causes (Residual)	3,124	17.4	54.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 45-54 years				
...	All causes	37,419	100.0	720.8
1	Malignant neoplasms (C00-C97)	8,800	23.5	169.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,612	23.0	165.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,856	7.6	55.0
4	Human immunodeficiency virus (HIV) disease (B20-B24)	2,449	6.5	47.2
5	Cerebrovascular diseases (I60-I69)	1,919	5.1	37.0
6	Diabetes mellitus (E10-E14)	1,418	3.8	27.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	841	2.2	16.2
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	825	2.2	15.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	804	2.1	15.5
10	Septicemia (A40-A41)	680	1.8	13.1
11	Chronic lower respiratory diseases (J40-J47)	621	1.7	12.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	570	1.5	11.0
13	Influenza and pneumonia (J10-J18)	457	1.2	8.8
14	Viral hepatitis (B15-B19)	396	1.1	7.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	310	0.8	6.0
...	All other causes (Residual)	5,861	15.7	112.9
Black, both sexes, 55-64 years				
...	All causes	46,396	100.0	1,440.3
1	Malignant neoplasms (C00-C97)	14,171	30.5	439.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,050	26.0	374.1
3	Cerebrovascular diseases (I60-I69)	2,613	5.6	81.1
4	Diabetes mellitus (E10-E14)	2,500	5.4	77.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,557	3.4	48.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,346	2.9	41.8
7	Chronic lower respiratory diseases (J40-J47)	1,205	2.6	37.4
8	Septicemia (A40-A41)	997	2.1	30.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	976	2.1	30.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	740	1.6	23.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	710	1.5	22.0
12	Influenza and pneumonia (J10-J18)	511	1.1	15.9
13	Viral hepatitis (B15-B19)	396	0.9	12.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	309	0.7	9.6
15	Aortic aneurysm and dissection (I71)	213	0.5	6.6
...	All other causes (Residual)	6,102	13.2	189.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 65-74 years				
...	All causes	51,126	100.0	2,835.9
1	Malignant neoplasms (C00-C97)	15,767	30.8	874.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,932	27.3	772.8
3	Cerebrovascular diseases (I60-I69)	3,209	6.3	178.0
4	Diabetes mellitus (E10-E14)	3,008	5.9	166.8
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,846	3.6	102.4
6	Chronic lower respiratory diseases (J40-J47)	1,818	3.6	100.8
7	Septicemia (A40-A41)	1,245	2.4	69.1
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	887	1.7	49.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	865	1.7	48.0
10	Influenza and pneumonia (J10-J18)	806	1.6	44.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	340	0.7	18.9
12	Alzheimer's disease (G30)	336	0.7	18.6
13	Aortic aneurysm and dissection (I71)	259	0.5	14.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	254	0.5	14.1
15	Pneumonitis due to solids and liquids (J69)	250	0.5	13.9
...	All other causes (Residual)	6,304	12.3	349.7
Black, both sexes, 75-84 years				
...	All causes	60,002	100.0	5,886.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,370	28.9	1,703.9
2	Malignant neoplasms (C00-C97)	14,237	23.7	1,396.6
3	Cerebrovascular diseases (I60-I69)	4,442	7.4	435.7
4	Diabetes mellitus (E10-E14)	3,181	5.3	312.0
5	Chronic lower respiratory diseases (J40-J47)	2,303	3.8	225.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,138	3.6	209.7
7	Septicemia (A40-A41)	1,532	2.6	150.3
8	Alzheimer's disease (G30)	1,495	2.5	146.7
9	Influenza and pneumonia (J10-J18)	1,425	2.4	139.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,193	2.0	117.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	824	1.4	80.8
12	Pneumonitis due to solids and liquids (J69)	429	0.7	42.1
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	322	0.5	31.6
14	Parkinson's disease (G20-G21)	306	0.5	30.0
15	Aortic aneurysm and dissection (I71)	285	0.5	28.0
...	All other causes (Residual)	8,520	14.2	835.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 85 years and over				
...	All causes	48,142	100.0	12,489.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,635	32.5	4,056.3
2	Malignant neoplasms (C00-C97)	6,469	13.4	1,678.3
3	Cerebrovascular diseases (I60-I69)	3,888	8.1	1,008.7
4	Alzheimer's disease (G30)	2,559	5.3	663.9
5	Diabetes mellitus (E10-E14)	1,816	3.8	471.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,706	3.5	442.6
7	Influenza and pneumonia (J10-J18)	1,675	3.5	434.6
8	Chronic lower respiratory diseases (J40-J47)	1,279	2.7	331.8
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,227	2.5	318.3
10	Septicemia (A40-A41)	1,108	2.3	287.5
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	580	1.2	150.5
12	Pneumonitis due to solids and liquids (J69)	495	1.0	128.4
13	Atherosclerosis (I70)	276	0.6	71.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	240	0.5	62.3
15	Parkinson's disease (G20-G21)	198	0.4	51.4
...	All other causes (Residual)	8,991	18.7	2,332.6
Black, male, all ages				
...	All causes	148,602	100.0	786.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,230	24.4	191.8
2	Malignant neoplasms (C00-C97)	32,556	21.9	172.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,605	6.5	50.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,677	5.2	40.6
5	Cerebrovascular diseases (I60-I69)	7,424	5.0	39.3
6	Diabetes mellitus (E10-E14)	5,772	3.9	30.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	4,443	3.0	23.5
8	Chronic lower respiratory diseases (J40-J47)	4,136	2.8	21.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,812	2.6	20.2
10	Certain conditions originating in the perinatal period (P00-P96)	2,811	1.9	14.9
11	Septicemia (A40-A41)	2,717	1.8	14.4
12	Influenza and pneumonia (J10-J18)	2,486	1.7	13.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,126	1.4	11.3
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,669	1.1	8.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,519	1.0	8.0
...	All other causes (Residual)	23,619	15.9	125.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 1-4 years				
...	All causes	633	100.0	47.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	214	33.8	15.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	84	13.3	6.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	42	6.6	3.1
4	Malignant neoplasms (C00-C97)	35	5.5	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	4.3	2.0
6	Influenza and pneumonia (J10-J18)	16	2.5	*
7	Septicemia (A40-A41)	12	1.9	*
7	Chronic lower respiratory diseases (J40-J47)	12	1.9	*
9	Anemias (D50-D64)	9	1.4	*
9	Certain conditions originating in the perinatal period (P00-P96)	9	1.4	*
11	Cerebrovascular diseases (I60-I69)	8	1.3	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.8	*
13	Acute bronchitis and bronchiolitis (J20-J21)	3	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
15	Meningococcal infection (A39)	1	0.2	*
15	Syphilis (A50-A53)	1	0.2	*
15	Meningitis (G00,G03)	1	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	149	23.5	11.1
Black, male, 5-14 years				
...	All causes	831	100.0	24.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	295	35.5	8.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	105	12.6	3.1
3	Malignant neoplasms (C00-C97)	83	10.0	2.5
4	Chronic lower respiratory diseases (J40-J47)	44	5.3	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	4.2	1.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	3.6	0.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	3.5	0.9
8	Septicemia (A40-A41)	13	1.6	*
9	Anemias (D50-D64)	9	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.8	*
10	Cerebrovascular diseases (I60-I69)	7	0.8	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.7	*
13	Meningitis (G00,G03)	3	0.4	*
13	Influenza and pneumonia (J10-J18)	3	0.4	*
13	Pneumonitis due to solids and liquids (J69)	3	0.4	*
...	All other causes (Residual)	159	19.1	4.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 15-24 years				
...	All causes	5,793	100.0	171.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,982	51.5	88.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,331	23.0	39.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	358	6.2	10.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	183	3.2	5.4
5	Malignant neoplasms (C00-C97)	153	2.6	4.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	84	1.5	2.5
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	53	0.9	1.6
8	Chronic lower respiratory diseases (J40-J47)	47	0.8	1.4
9	Anemias (D50-D64)	37	0.6	1.1
10	Diabetes mellitus (E10-E14)	32	0.6	0.9
11	Legal intervention (Y35,Y89.0)	26	0.4	0.8
12	Influenza and pneumonia (J10-J18)	23	0.4	0.7
13	Septicemia (A40-A41)	21	0.4	0.6
14	Cerebrovascular diseases (I60-I69)	20	0.3	0.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.2	*
...	All other causes (Residual)	421	7.3	12.5
Black, male, 25-34 years				
...	All causes	6,919	100.0	254.2
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,346	33.9	86.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,503	21.7	55.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	663	9.6	24.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	421	6.1	15.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	418	6.0	15.4
6	Malignant neoplasms (C00-C97)	272	3.9	10.0
7	Diabetes mellitus (E10-E14)	137	2.0	5.0
8	Cerebrovascular diseases (I60-I69)	75	1.1	2.8
9	Chronic lower respiratory diseases (J40-J47)	61	0.9	2.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	57	0.8	2.1
11	Anemias (D50-D64)	55	0.8	2.0
12	Septicemia (A40-A41)	45	0.7	1.7
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	44	0.6	1.6
13	Legal intervention (Y35,Y89.0)	44	0.6	1.6
15	Influenza and pneumonia (J10-J18)	37	0.5	1.4
...	All other causes (Residual)	741	10.7	27.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 35-44 years				
...	All causes	10,521	100.0	392.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,892	18.0	70.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,669	15.9	62.2
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,358	12.9	50.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,068	10.2	39.8
5	Malignant neoplasms (C00-C97)	980	9.3	36.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	352	3.3	13.1
7	Diabetes mellitus (E10-E14)	346	3.3	12.9
8	Cerebrovascular diseases (I60-I69)	342	3.3	12.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	183	1.7	6.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	155	1.5	5.8
11	Septicemia (A40-A41)	134	1.3	5.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	127	1.2	4.7
13	Chronic lower respiratory diseases (J40-J47)	103	1.0	3.8
14	Influenza and pneumonia (J10-J18)	98	0.9	3.7
15	Anemias (D50-D64)	65	0.6	2.4
...	All other causes (Residual)	1,649	15.7	61.5
Black, male, 45-54 years				
...	All causes	22,112	100.0	921.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,511	24.9	229.8
2	Malignant neoplasms (C00-C97)	4,369	19.8	182.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,055	9.3	85.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,646	7.4	68.6
5	Cerebrovascular diseases (I60-I69)	1,044	4.7	43.5
6	Diabetes mellitus (E10-E14)	804	3.6	33.5
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	650	2.9	27.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	542	2.5	22.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	480	2.2	20.0
10	Septicemia (A40-A41)	354	1.6	14.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	332	1.5	13.8
12	Chronic lower respiratory diseases (J40-J47)	319	1.4	13.3
13	Influenza and pneumonia (J10-J18)	276	1.2	11.5
14	Viral hepatitis (B15-B19)	259	1.2	10.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	251	1.1	10.5
...	All other causes (Residual)	3,220	14.6	134.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 55-64 years				
...	All causes	27,198	100.0	1,891.8
1	Malignant neoplasms (C00-C97)	7,805	28.7	542.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,568	27.8	526.4
3	Cerebrovascular diseases (I60-I69)	1,523	5.6	105.9
4	Diabetes mellitus (E10-E14)	1,306	4.8	90.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,122	4.1	78.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	717	2.6	49.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	709	2.6	49.3
8	Chronic lower respiratory diseases (J40-J47)	690	2.5	48.0
9	Septicemia (A40-A41)	534	2.0	37.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	481	1.8	33.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	404	1.5	28.1
12	Influenza and pneumonia (J10-J18)	306	1.1	21.3
13	Viral hepatitis (B15-B19)	283	1.0	19.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	245	0.9	17.0
15	Aortic aneurysm and dissection (I71)	138	0.5	9.6
...	All other causes (Residual)	3,367	12.4	234.2
Black, male, 65-74 years				
...	All causes	27,587	100.0	3,669.2
1	Malignant neoplasms (C00-C97)	8,695	31.5	1,156.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,854	28.5	1,044.6
3	Cerebrovascular diseases (I60-I69)	1,644	6.0	218.7
4	Diabetes mellitus (E10-E14)	1,415	5.1	188.2
5	Chronic lower respiratory diseases (J40-J47)	1,028	3.7	136.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	863	3.1	114.8
7	Septicemia (A40-A41)	570	2.1	75.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	548	2.0	72.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	450	1.6	59.9
10	Influenza and pneumonia (J10-J18)	438	1.6	58.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	220	0.8	29.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	179	0.6	23.8
13	Pneumonitis due to solids and liquids (J69)	155	0.6	20.6
14	Alzheimer's disease (G30)	143	0.5	19.0
14	Aortic aneurysm and dissection (I71)	143	0.5	19.0
...	All other causes (Residual)	3,242	11.8	431.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 75-84 years				
...	All causes	27,322	100.0	7,393.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,871	28.8	2,129.9
2	Malignant neoplasms (C00-C97)	7,314	26.8	1,979.1
3	Cerebrovascular diseases (I60-I69)	1,741	6.4	471.1
4	Chronic lower respiratory diseases (J40-J47)	1,253	4.6	339.1
5	Diabetes mellitus (E10-E14)	1,211	4.4	327.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	927	3.4	250.8
7	Influenza and pneumonia (J10-J18)	669	2.4	181.0
8	Septicemia (A40-A41)	648	2.4	175.3
9	Alzheimer's disease (G30)	491	1.8	132.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	467	1.7	126.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	428	1.6	115.8
12	Pneumonitis due to solids and liquids (J69)	247	0.9	66.8
13	Parkinson's disease (G20-G21)	187	0.7	50.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	153	0.6	41.4
15	Aortic aneurysm and dissection (I71)	110	0.4	29.8
...	All other causes (Residual)	3,605	13.2	975.5
Black, male, 85 years and over				
...	All causes	14,778	100.0	13,206.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,558	30.8	4,073.1
2	Malignant neoplasms (C00-C97)	2,846	19.3	2,543.3
3	Cerebrovascular diseases (I60-I69)	987	6.7	882.0
4	Chronic lower respiratory diseases (J40-J47)	573	3.9	512.0
5	Influenza and pneumonia (J10-J18)	565	3.8	504.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	547	3.7	488.8
7	Alzheimer's disease (G30)	527	3.6	470.9
8	Diabetes mellitus (E10-E14)	519	3.5	463.8
9	Septicemia (A40-A41)	332	2.2	296.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	304	2.1	271.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	223	1.5	199.3
12	Pneumonitis due to solids and liquids (J69)	194	1.3	173.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	90	0.6	80.4
14	Atherosclerosis (I70)	84	0.6	75.1
15	Parkinson's disease (G20-G21)	79	0.5	70.6
...	All other causes (Residual)	2,350	15.9	2,100.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, all ages				
...	All causes	141,369	100.0	684.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,023	25.5	174.3
2	Malignant neoplasms (C00-C97)	30,526	21.6	147.7
3	Cerebrovascular diseases (I60-I69)	9,621	6.8	46.5
4	Diabetes mellitus (E10-E14)	7,041	5.0	34.1
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,585	3.2	22.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,312	3.1	20.9
7	Chronic lower respiratory diseases (J40-J47)	3,594	2.5	17.4
8	Septicemia (A40-A41)	3,391	2.4	16.4
9	Alzheimer's disease (G30)	3,265	2.3	15.8
10	Influenza and pneumonia (J10-J18)	2,825	2.0	13.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,774	2.0	13.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,411	1.7	11.7
13	Certain conditions originating in the perinatal period (P00-P96)	2,310	1.6	11.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,355	1.0	6.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	861	0.6	4.2
...	All other causes (Residual)	26,475	18.7	128.1
Black, female, 1-4 years				
...	All causes	511	100.0	39.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	157	30.7	12.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	13.7	5.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	12.7	5.0
4	Malignant neoplasms (C00-C97)	30	5.9	2.3
5	Influenza and pneumonia (J10-J18)	18	3.5	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	3.3	*
7	Certain conditions originating in the perinatal period (P00-P96)	13	2.5	*
8	Septicemia (A40-A41)	9	1.8	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.8	*
9	Cerebrovascular diseases (I60-I69)	4	0.8	*
11	Anemias (D50-D64)	3	0.6	*
11	Meningitis (G00,G03)	3	0.6	*
11	Chronic lower respiratory diseases (J40-J47)	3	0.6	*
14	Diabetes mellitus (E10-E14)	2	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
...	All other causes (Residual)	111	21.7	8.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 5-14 years				
...	All causes	563	100.0	17.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	176	31.3	5.4
2	Malignant neoplasms (C00-C97)	62	11.0	1.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	7.8	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	6.7	1.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	5.7	1.0
6	Chronic lower respiratory diseases (J40-J47)	25	4.4	0.8
7	Cerebrovascular diseases (I60-I69)	14	2.5	*
8	Anemias (D50-D64)	11	2.0	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.4	*
10	Septicemia (A40-A41)	7	1.2	*
10	Diabetes mellitus (E10-E14)	7	1.2	*
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	1.2	*
13	Influenza and pneumonia (J10-J18)	6	1.1	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.9	*
14	Meningitis (G00,G03)	5	0.9	*
...	All other causes (Residual)	116	20.6	3.6
Black, female, 15-24 years				
...	All causes	1,690	100.0	51.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	419	24.8	12.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	313	18.5	9.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	125	7.4	3.8
4	Malignant neoplasms (C00-C97)	111	6.6	3.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	60	3.6	1.8
6	Pregnancy, childbirth and the puerperium (O00-099)	58	3.4	1.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	58	3.4	1.8
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	2.1	1.1
9	Chronic lower respiratory diseases (J40-J47)	28	1.7	0.9
10	Anemias (D50-D64)	27	1.6	0.8
11	Cerebrovascular diseases (I60-I69)	24	1.4	0.7
12	Diabetes mellitus (E10-E14)	23	1.4	0.7
13	Septicemia (A40-A41)	20	1.2	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	1.1	*
15	Influenza and pneumonia (J10-J18)	18	1.1	*
...	All other causes (Residual)	352	20.8	10.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 25-34 years				
...	All causes	3,126	100.0	106.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	430	13.8	14.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	370	11.8	12.6
3	Malignant neoplasms (C00-C97)	367	11.7	12.5
4	Human immunodeficiency virus (HIV) disease (B20-B24)	352	11.3	12.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	324	10.4	11.1
6	Pregnancy, childbirth and the puerperium (O00-099)	118	3.8	4.0
7	Diabetes mellitus (E10-E14)	103	3.3	3.5
8	Cerebrovascular diseases (I60-I69)	76	2.4	2.6
9	Anemias (D50-D64)	57	1.8	1.9
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	53	1.7	1.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	1.6	1.7
12	Chronic lower respiratory diseases (J40-J47)	47	1.5	1.6
13	Septicemia (A40-A41)	44	1.4	1.5
14	Influenza and pneumonia (J10-J18)	40	1.3	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	0.9	1.0
...	All other causes (Residual)	666	21.3	22.7
Black, female, 35-44 years				
...	All causes	7,409	100.0	245.0
1	Malignant neoplasms (C00-C97)	1,535	20.7	50.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,186	16.0	39.2
3	Human immunodeficiency virus (HIV) disease (B20-B24)	831	11.2	27.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	678	9.2	22.4
5	Cerebrovascular diseases (I60-I69)	355	4.8	11.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	279	3.8	9.2
7	Diabetes mellitus (E10-E14)	237	3.2	7.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	153	2.1	5.1
9	Septicemia (A40-A41)	139	1.9	4.6
10	Chronic lower respiratory diseases (J40-J47)	123	1.7	4.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	101	1.4	3.3
12	Influenza and pneumonia (J10-J18)	88	1.2	2.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	82	1.1	2.7
14	Anemias (D50-D64)	81	1.1	2.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	66	0.9	2.2
...	All other causes (Residual)	1,475	19.9	48.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 45-54 years				
...	All causes	15,307	100.0	548.1
1	Malignant neoplasms (C00-C97)	4,431	28.9	158.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,101	20.3	111.0
3	Cerebrovascular diseases (I60-I69)	875	5.7	31.3
4	Human immunodeficiency virus (HIV) disease (B20-B24)	803	5.2	28.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	801	5.2	28.7
6	Diabetes mellitus (E10-E14)	614	4.0	22.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	361	2.4	12.9
8	Septicemia (A40-A41)	326	2.1	11.7
9	Chronic lower respiratory diseases (J40-J47)	302	2.0	10.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	262	1.7	9.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	238	1.6	8.5
12	Influenza and pneumonia (J10-J18)	181	1.2	6.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	175	1.1	6.3
14	Viral hepatitis (B15-B19)	137	0.9	4.9
15	Anemias (D50-D64)	76	0.5	2.7
...	All other causes (Residual)	2,624	17.1	94.0
Black, female, 55-64 years				
...	All causes	19,198	100.0	1,076.3
1	Malignant neoplasms (C00-C97)	6,366	33.2	356.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,482	23.3	251.3
3	Diabetes mellitus (E10-E14)	1,194	6.2	66.9
4	Cerebrovascular diseases (I60-I69)	1,090	5.7	61.1
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	629	3.3	35.3
6	Chronic lower respiratory diseases (J40-J47)	515	2.7	28.9
7	Septicemia (A40-A41)	463	2.4	26.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	435	2.3	24.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	336	1.8	18.8
10	Human immunodeficiency virus (HIV) disease (B20-B24)	267	1.4	15.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	229	1.2	12.8
12	Influenza and pneumonia (J10-J18)	205	1.1	11.5
13	Viral hepatitis (B15-B19)	113	0.6	6.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	84	0.4	4.7
15	Aortic aneurysm and dissection (I71)	75	0.4	4.2
...	All other causes (Residual)	2,715	14.1	152.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 65-74 years				
...	All causes	23,539	100.0	2,239.7
1	Malignant neoplasms (C00-C97)	7,072	30.0	672.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,078	25.8	578.3
3	Diabetes mellitus (E10-E14)	1,593	6.8	151.6
4	Cerebrovascular diseases (I60-I69)	1,565	6.6	148.9
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	983	4.2	93.5
6	Chronic lower respiratory diseases (J40-J47)	790	3.4	75.2
7	Septicemia (A40-A41)	675	2.9	64.2
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	437	1.9	41.6
9	Influenza and pneumonia (J10-J18)	368	1.6	35.0
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	317	1.3	30.2
11	Alzheimer's disease (G30)	193	0.8	18.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	120	0.5	11.4
13	Aortic aneurysm and dissection (I71)	116	0.5	11.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	99	0.4	9.4
15	Pneumonitis due to solids and liquids (J69)	95	0.4	9.0
...	All other causes (Residual)	3,038	12.9	289.1
Black, female, 75-84 years				
...	All causes	32,680	100.0	5,028.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,499	29.1	1,461.7
2	Malignant neoplasms (C00-C97)	6,923	21.2	1,065.3
3	Cerebrovascular diseases (I60-I69)	2,701	8.3	415.6
4	Diabetes mellitus (E10-E14)	1,970	6.0	303.2
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,211	3.7	186.4
6	Chronic lower respiratory diseases (J40-J47)	1,050	3.2	161.6
7	Alzheimer's disease (G30)	1,004	3.1	154.5
8	Septicemia (A40-A41)	884	2.7	136.0
9	Influenza and pneumonia (J10-J18)	756	2.3	116.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	726	2.2	111.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	396	1.2	60.9
12	Pneumonitis due to solids and liquids (J69)	182	0.6	28.0
13	Aortic aneurysm and dissection (I71)	175	0.5	26.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	169	0.5	26.0
15	Parkinson's disease (G20-G21)	119	0.4	18.3
...	All other causes (Residual)	4,915	15.0	756.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 85 years and over				
...	All causes	33,364	100.0	12,196.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,077	33.2	4,049.4
2	Malignant neoplasms (C00-C97)	3,623	10.9	1,324.4
3	Cerebrovascular diseases (I60-I69)	2,901	8.7	1,060.5
4	Alzheimer's disease (G30)	2,032	6.1	742.8
5	Diabetes mellitus (E10-E14)	1,297	3.9	474.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,159	3.5	423.7
7	Influenza and pneumonia (J10-J18)	1,110	3.3	405.8
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	923	2.8	337.4
9	Septicemia (A40-A41)	776	2.3	283.7
10	Chronic lower respiratory diseases (J40-J47)	706	2.1	258.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	357	1.1	130.5
12	Pneumonitis due to solids and liquids (J69)	301	0.9	110.0
13	Atherosclerosis (I70)	192	0.6	70.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	150	0.4	54.8
15	Aortic aneurysm and dissection (I71)	125	0.4	45.7
...	All other causes (Residual)	6,635	19.9	2,425.5
American Indian, both sexes, all ages				
...	All causes	14,037	100.0	438.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,736	19.5	85.5
2	Malignant neoplasms (C00-C97)	2,447	17.4	76.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,704	12.1	53.2
4	Diabetes mellitus (E10-E14)	811	5.8	25.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	596	4.2	18.6
6	Cerebrovascular diseases (I60-I69)	548	3.9	17.1
7	Chronic lower respiratory diseases (J40-J47)	508	3.6	15.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	395	2.8	12.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	288	2.1	9.0
10	Influenza and pneumonia (J10-J18)	267	1.9	8.3
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	254	1.8	7.9
12	Septicemia (A40-A41)	207	1.5	6.5
13	Alzheimer's disease (G30)	169	1.2	5.3
14	Certain conditions originating in the perinatal period (P00-P96)	137	1.0	4.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	117	0.8	3.7
...	All other causes (Residual)	2,853	20.3	89.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 1-4 years				
...	All causes	94	100.0	54.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32	34.0	18.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	11.7	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	10.6	*
4	Malignant neoplasms (C00-C97)	7	7.4	*
5	Influenza and pneumonia (J10-J18)	4	4.3	*
6	Septicemia (A40-A41)	2	2.1	*
6	Cerebrovascular diseases (I60-I69)	2	2.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	1.1	*
...	All other causes (Residual)	25	26.6	14.5
American Indian, both sexes, 5-14 years				
...	All causes	91	100.0	16.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	43	47.3	8.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	13.2	*
3	Malignant neoplasms (C00-C97)	5	5.5	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	5.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	3.3	*
6	Cerebrovascular diseases (I60-I69)	2	2.2	*
6	Influenza and pneumonia (J10-J18)	2	2.2	*
6	Chronic lower respiratory diseases (J40-J47)	2	2.2	*
6	Diseases of appendix (K35-K38)	2	2.2	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	2.2	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.1	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.1	*
...	All other causes (Residual)	11	12.1	*

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 15-24 years				
...	All causes	664	100.0	110.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	328	49.4	54.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	136	20.5	22.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	77	11.6	12.9
4	Malignant neoplasms (C00-C97)	16	2.4	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	2.1	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	1.7	*
7	Septicemia (A40-A41)	3	0.5	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
7	Cerebrovascular diseases (I60-I69)	3	0.5	*
7	Influenza and pneumonia (J10-J18)	3	0.5	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.5	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
7	Legal intervention (Y35,Y89.0)	3	0.5	*
14	Pregnancy, childbirth and the puerperium (O00-099)	2	0.3	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
15	Diabetes mellitus (E10-E14)	1	0.2	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
15	Aortic aneurysm and dissection (I71)	1	0.2	*
15	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
...	All other causes (Residual)	53	8.0	8.8
American Indian, both sexes, 25-34 years				
...	All causes	708	100.0	145.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	305	43.1	62.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	77	10.9	15.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	9.5	13.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	4.9	7.2
5	Malignant neoplasms (C00-C97)	32	4.5	6.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	31	4.4	6.4
7	Diabetes mellitus (E10-E14)	15	2.1	*
8	Influenza and pneumonia (J10-J18)	7	1.0	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.0	*
10	Septicemia (A40-A41)	6	0.8	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.7	*
11	Cerebrovascular diseases (I60-I69)	5	0.7	*
13	Pregnancy, childbirth and the puerperium (O00-099)	4	0.6	*
14	Legal intervention (Y35,Y89.0)	3	0.4	*
15	Tuberculosis (A16-A19)	2	0.3	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.3	*
...	All other causes (Residual)	105	14.8	21.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 35-44 years				
...	All causes	1,252	100.0	272.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	300	24.0	65.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	141	11.3	30.6
3	Chronic liver disease and cirrhosis (K70,K73-K74)	121	9.7	26.3
4	Malignant neoplasms (C00-C97)	93	7.4	20.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	86	6.9	18.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	3.8	10.4
7	Diabetes mellitus (E10-E14)	39	3.1	8.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	34	2.7	7.4
9	Cerebrovascular diseases (I60-I69)	31	2.5	6.7
10	Viral hepatitis (B15-B19)	18	1.4	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	1.4	*
12	Influenza and pneumonia (J10-J18)	16	1.3	*
13	Septicemia (A40-A41)	13	1.0	*
14	Chronic lower respiratory diseases (J40-J47)	12	1.0	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.5	*
...	All other causes (Residual)	277	22.1	60.2
American Indian, both sexes, 45-54 years				
...	All causes	1,937	100.0	463.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	332	17.1	79.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	313	16.2	74.9
3	Malignant neoplasms (C00-C97)	308	15.9	73.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	201	10.4	48.1
5	Diabetes mellitus (E10-E14)	103	5.3	24.6
6	Cerebrovascular diseases (I60-I69)	52	2.7	12.4
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	51	2.6	12.2
8	Viral hepatitis (B15-B19)	41	2.1	9.8
9	Septicemia (A40-A41)	37	1.9	8.8
10	Chronic lower respiratory diseases (J40-J47)	33	1.7	7.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	1.7	7.9
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	1.5	7.2
13	Influenza and pneumonia (J10-J18)	24	1.2	5.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	23	1.2	5.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	0.5	*
...	All other causes (Residual)	346	17.9	82.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 55-64 years				
...	All causes	2,226	100.0	851.5
1	Malignant neoplasms (C00-C97)	543	24.4	207.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	492	22.1	188.2
3	Diabetes mellitus (E10-E14)	190	8.5	72.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	157	7.1	60.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	135	6.1	51.6
6	Chronic lower respiratory diseases (J40-J47)	86	3.9	32.9
7	Cerebrovascular diseases (I60-I69)	66	3.0	25.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	52	2.3	19.9
9	Septicemia (A40-A41)	40	1.8	15.3
10	Viral hepatitis (B15-B19)	28	1.3	10.7
11	Influenza and pneumonia (J10-J18)	26	1.2	9.9
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	1.0	8.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	0.6	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	0.4	*
15	Pneumonitis due to solids and liquids (J69)	7	0.3	*
...	All other causes (Residual)	359	16.1	137.3
American Indian, both sexes, 65-74 years				
...	All causes	2,452	100.0	1,884.2
1	Malignant neoplasms (C00-C97)	680	27.7	522.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	571	23.3	438.8
3	Diabetes mellitus (E10-E14)	190	7.7	146.0
4	Chronic lower respiratory diseases (J40-J47)	137	5.6	105.3
5	Cerebrovascular diseases (I60-I69)	105	4.3	80.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	85	3.5	65.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	81	3.3	62.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	73	3.0	56.1
9	Influenza and pneumonia (J10-J18)	40	1.6	30.7
10	Septicemia (A40-A41)	37	1.5	28.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	22	0.9	16.9
12	Alzheimer's disease (G30)	15	0.6	*
12	Pneumonitis due to solids and liquids (J69)	15	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.4	*
14	Parkinson's disease (G20-G21)	11	0.4	*
14	Aortic aneurysm and dissection (I71)	11	0.4	*
...	All other causes (Residual)	368	15.0	282.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 75-84 years				
...	All causes	2,542	100.0	3,939.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	671	26.4	1,039.8
2	Malignant neoplasms (C00-C97)	560	22.0	867.8
3	Diabetes mellitus (E10-E14)	184	7.2	285.1
4	Chronic lower respiratory diseases (J40-J47)	166	6.5	257.2
5	Cerebrovascular diseases (I60-I69)	147	5.8	227.8
6	Alzheimer's disease (G30)	73	2.9	113.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	68	2.7	105.4
8	Influenza and pneumonia (J10-J18)	67	2.6	103.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	52	2.0	80.6
10	Septicemia (A40-A41)	45	1.8	69.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	25	1.0	38.7
12	Parkinson's disease (G20-G21)	24	0.9	37.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	23	0.9	35.6
14	Pneumonitis due to solids and liquids (J69)	16	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.6	*
...	All other causes (Residual)	406	16.0	629.1
American Indian, both sexes, 85 years and over				
...	All causes	1,676	100.0	6,927.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	470	28.0	1,942.6
2	Malignant neoplasms (C00-C97)	203	12.1	839.1
3	Cerebrovascular diseases (I60-I69)	133	7.9	549.7
4	Diabetes mellitus (E10-E14)	89	5.3	367.9
5	Alzheimer's disease (G30)	80	4.8	330.7
6	Influenza and pneumonia (J10-J18)	75	4.5	310.0
7	Chronic lower respiratory diseases (J40-J47)	68	4.1	281.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	2.4	169.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36	2.1	148.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	33	2.0	136.4
11	Pneumonitis due to solids and liquids (J69)	23	1.4	95.1
12	Septicemia (A40-A41)	21	1.3	86.8
13	Parkinson's disease (G20-G21)	20	1.2	82.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.1	*
15	Atherosclerosis (I70)	11	0.7	*
...	All other causes (Residual)	355	21.2	1,467.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, all ages				
...	All causes	7,630	100.0	477.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,532	20.1	95.8
2	Malignant neoplasms (C00-C97)	1,217	16.0	76.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,184	15.5	74.0
4	Diabetes mellitus (E10-E14)	362	4.7	22.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	330	4.3	20.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	309	4.0	19.3
7	Chronic lower respiratory diseases (J40-J47)	234	3.1	14.6
8	Cerebrovascular diseases (I60-I69)	231	3.0	14.4
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	206	2.7	12.9
10	Influenza and pneumonia (J10-J18)	137	1.8	8.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	135	1.8	8.4
12	Septicemia (A40-A41)	99	1.3	6.2
13	Certain conditions originating in the perinatal period (P00-P96)	91	1.2	5.7
14	Viral hepatitis (B15-B19)	66	0.9	4.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	0.9	4.1
...	All other causes (Residual)	1,432	18.8	89.6
American Indian, male, 1-4 years				
...	All causes	51	100.0	58.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	29.4	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	17.6	*
3	Influenza and pneumonia (J10-J18)	3	5.9	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	5.9	*
5	Malignant neoplasms (C00-C97)	2	3.9	*
5	Cerebrovascular diseases (I60-I69)	2	3.9	*
7	Certain conditions originating in the perinatal period (P00-P96)	1	2.0	*
...	All other causes (Residual)	16	31.4	*
American Indian, male, 5-14 years				
...	All causes	47	100.0	17.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	55.3	9.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	8.5	*
3	Malignant neoplasms (C00-C97)	3	6.4	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	4.3	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	4.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.1	*
6	Cerebrovascular diseases (I60-I69)	1	2.1	*
6	Chronic lower respiratory diseases (J40-J47)	1	2.1	*
6	Diseases of appendix (K35-K38)	1	2.1	*
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	2.1	*
...	All other causes (Residual)	5	10.6	*

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 15-24 years				
...	All causes	478	100.0	156.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	237	49.6	77.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	110	23.0	35.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	14.4	22.5
4	Malignant neoplasms (C00-C97)	10	2.1	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	1.9	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	1.0	*
7	Legal intervention (Y35,Y89.0)	3	0.6	*
8	Septicemia (A40-A41)	2	0.4	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
10	Diabetes mellitus (E10-E14)	1	0.2	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
10	Cerebrovascular diseases (I60-I69)	1	0.2	*
10	Influenza and pneumonia (J10-J18)	1	0.2	*
10	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
...	All other causes (Residual)	25	5.2	8.2
American Indian, male, 25-34 years				
...	All causes	492	100.0	194.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	225	45.7	88.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	63	12.8	24.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	10.8	20.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	4.7	9.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	21	4.3	8.3
6	Malignant neoplasms (C00-C97)	13	2.6	*
7	Diabetes mellitus (E10-E14)	6	1.2	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	5	1.0	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.8	*
10	Cerebrovascular diseases (I60-I69)	3	0.6	*
10	Legal intervention (Y35,Y89.0)	3	0.6	*
12	Influenza and pneumonia (J10-J18)	2	0.4	*
13	Tuberculosis (A16-A19)	1	0.2	*
13	Viral hepatitis (B15-B19)	1	0.2	*
13	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
13	Pneumonitis due to solids and liquids (J69)	1	0.2	*
13	Peptic ulcer (K25-K28)	1	0.2	*
...	All other causes (Residual)	66	13.4	26.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 35-44 years				
...	All causes	784	100.0	338.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	210	26.8	90.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	104	13.3	44.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	63	8.0	27.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	59	7.5	25.5
5	Malignant neoplasms (C00-C97)	35	4.5	15.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	34	4.3	14.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	21	2.7	9.1
7	Cerebrovascular diseases (I60-I69)	21	2.7	9.1
9	Diabetes mellitus (E10-E14)	19	2.4	*
10	Viral hepatitis (B15-B19)	16	2.0	*
11	Influenza and pneumonia (J10-J18)	10	1.3	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.3	*
13	Chronic lower respiratory diseases (J40-J47)	7	0.9	*
14	Septicemia (A40-A41)	5	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.5	*
...	All other causes (Residual)	166	21.2	71.7
American Indian, male, 45-54 years				
...	All causes	1,199	100.0	591.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	242	20.2	119.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	229	19.1	113.1
3	Malignant neoplasms (C00-C97)	151	12.6	74.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	119	9.9	58.7
5	Diabetes mellitus (E10-E14)	56	4.7	27.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	3.3	19.3
7	Cerebrovascular diseases (I60-I69)	33	2.8	16.3
8	Viral hepatitis (B15-B19)	30	2.5	14.8
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	1.8	10.4
10	Septicemia (A40-A41)	19	1.6	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	18	1.5	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	1.5	*
13	Chronic lower respiratory diseases (J40-J47)	15	1.3	*
14	Influenza and pneumonia (J10-J18)	14	1.2	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	0.8	*
...	All other causes (Residual)	186	15.5	91.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 55-64 years				
...	All causes	1,294	100.0	1,029.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	322	24.9	256.2
2	Malignant neoplasms (C00-C97)	280	21.6	222.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	114	8.8	90.7
4	Diabetes mellitus (E10-E14)	103	8.0	81.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	82	6.3	65.2
6	Cerebrovascular diseases (I60-I69)	44	3.4	35.0
6	Chronic lower respiratory diseases (J40-J47)	44	3.4	35.0
8	Septicemia (A40-A41)	23	1.8	18.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	1.6	16.7
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	1.5	15.9
11	Viral hepatitis (B15-B19)	15	1.2	*
12	Influenza and pneumonia (J10-J18)	11	0.9	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	0.7	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.3	*
15	Pneumonitis due to solids and liquids (J69)	4	0.3	*
...	All other causes (Residual)	191	14.8	152.0
American Indian, male, 65-74 years				
...	All causes	1,295	100.0	2,146.7
1	Malignant neoplasms (C00-C97)	352	27.2	583.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	346	26.7	573.6
3	Diabetes mellitus (E10-E14)	82	6.3	135.9
4	Chronic lower respiratory diseases (J40-J47)	62	4.8	102.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	55	4.2	91.2
6	Cerebrovascular diseases (I60-I69)	50	3.9	82.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	37	2.9	61.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36	2.8	59.7
9	Influenza and pneumonia (J10-J18)	27	2.1	44.8
10	Septicemia (A40-A41)	21	1.6	34.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	1.0	*
12	Alzheimer's disease (G30)	9	0.7	*
12	Pneumonitis due to solids and liquids (J69)	9	0.7	*
14	Parkinson's disease (G20-G21)	7	0.5	*
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	7	0.5	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	0.5	*
...	All other causes (Residual)	175	13.5	290.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 75-84 years				
...	All causes	1,156	100.0	4,198.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	324	28.0	1,176.6
2	Malignant neoplasms (C00-C97)	280	24.2	1,016.8
3	Chronic lower respiratory diseases (J40-J47)	73	6.3	265.1
4	Diabetes mellitus (E10-E14)	66	5.7	239.7
5	Cerebrovascular diseases (I60-I69)	48	4.2	174.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39	3.4	141.6
7	Influenza and pneumonia (J10-J18)	36	3.1	130.7
8	Alzheimer's disease (G30)	25	2.2	90.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	2.1	87.2
10	Septicemia (A40-A41)	19	1.6	*
11	Parkinson's disease (G20-G21)	17	1.5	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11	1.0	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	10	0.9	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.7	*
14	Aortic aneurysm and dissection (I71)	8	0.7	*
14	Pneumonitis due to solids and liquids (J69)	8	0.7	*
...	All other causes (Residual)	160	13.8	581.0
American Indian, male, 85 years and over				
...	All causes	591	100.0	7,540.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	162	27.4	2,066.9
2	Malignant neoplasms (C00-C97)	91	15.4	1,161.0
3	Influenza and pneumonia (J10-J18)	33	5.6	421.0
4	Diabetes mellitus (E10-E14)	29	4.9	370.0
5	Alzheimer's disease (G30)	28	4.7	357.2
5	Chronic lower respiratory diseases (J40-J47)	28	4.7	357.2
7	Cerebrovascular diseases (I60-I69)	27	4.6	344.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	3.0	*
9	Pneumonitis due to solids and liquids (J69)	13	2.2	*
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13	2.2	*
11	Septicemia (A40-A41)	9	1.5	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	1.5	*
13	Parkinson's disease (G20-G21)	8	1.4	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.2	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	3	0.5	*
...	All other causes (Residual)	113	19.1	1,441.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, all ages				
...	All causes	6,407	100.0	399.9
1	Malignant neoplasms (C00-C97)	1,230	19.2	76.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,204	18.8	75.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	520	8.1	32.5
4	Diabetes mellitus (E10-E14)	449	7.0	28.0
5	Cerebrovascular diseases (I60-I69)	317	4.9	19.8
6	Chronic lower respiratory diseases (J40-J47)	274	4.3	17.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	266	4.2	16.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	153	2.4	9.5
9	Influenza and pneumonia (J10-J18)	130	2.0	8.1
10	Septicemia (A40-A41)	108	1.7	6.7
11	Alzheimer's disease (G30)	107	1.7	6.7
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	86	1.3	5.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	54	0.8	3.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	0.8	3.2
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	0.7	3.0
...	All other causes (Residual)	1,409	22.0	87.9
American Indian, female, 1-4 years				
...	All causes	43	100.0	50.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	39.5	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	16.3	*
3	Malignant neoplasms (C00-C97)	5	11.6	*
4	Septicemia (A40-A41)	2	4.7	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	4.7	*
6	Influenza and pneumonia (J10-J18)	1	2.3	*
...	All other causes (Residual)	9	20.9	*
American Indian, female, 5-14 years				
...	All causes	44	100.0	16.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	38.6	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	18.2	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	6.8	*
4	Malignant neoplasms (C00-C97)	2	4.5	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	4.5	*
4	Influenza and pneumonia (J10-J18)	2	4.5	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	2.3	*
7	Cerebrovascular diseases (I60-I69)	1	2.3	*
7	Chronic lower respiratory diseases (J40-J47)	1	2.3	*
7	Diseases of appendix (K35-K38)	1	2.3	*
...	All other causes (Residual)	6	13.6	*

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 15-24 years				
...	All causes	186	100.0	63.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	91	48.9	31.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26	14.0	8.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	4.8	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	4.3	*
5	Malignant neoplasms (C00-C97)	6	3.2	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.6	*
7	Cerebrovascular diseases (I60-I69)	2	1.1	*
7	Influenza and pneumonia (J10-J18)	2	1.1	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2	1.1	*
7	Pregnancy, childbirth and the puerperium (O00-099)	2	1.1	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.1	*
12	Septicemia (A40-A41)	1	0.5	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.5	*
12	Aortic aneurysm and dissection (I71)	1	0.5	*
12	Pneumonitis due to solids and liquids (J69)	1	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	*
...	All other causes (Residual)	28	15.1	9.6
American Indian, female, 25-34 years				
...	All causes	216	100.0	92.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80	37.0	34.1
2	Malignant neoplasms (C00-C97)	19	8.8	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	6.5	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	6.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	5.6	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	10	4.6	*
7	Diabetes mellitus (E10-E14)	9	4.2	*
8	Septicemia (A40-A41)	6	2.8	*
9	Influenza and pneumonia (J10-J18)	5	2.3	*
10	Pregnancy, childbirth and the puerperium (O00-099)	4	1.9	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.4	*
12	Cerebrovascular diseases (I60-I69)	2	0.9	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.9	*
14	Tuberculosis (A16-A19)	1	0.5	*
14	Anemias (D50-D64)	1	0.5	*
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.5	*
...	All other causes (Residual)	33	15.3	14.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 35-44 years				
...	All causes	468	100.0	204.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	90	19.2	39.4
2	Chronic liver disease and cirrhosis (K70,K73-K74)	62	13.2	27.1
3	Malignant neoplasms (C00-C97)	58	12.4	25.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	7.9	16.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23	4.9	10.1
6	Diabetes mellitus (E10-E14)	20	4.3	8.7
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	3.0	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	13	2.8	*
9	Cerebrovascular diseases (I60-I69)	10	2.1	*
10	Septicemia (A40-A41)	8	1.7	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.5	*
12	Influenza and pneumonia (J10-J18)	6	1.3	*
13	Chronic lower respiratory diseases (J40-J47)	5	1.1	*
14	Pregnancy, childbirth and the puerperium (O00-099)	3	0.6	*
15	Viral hepatitis (B15-B19)	2	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
15	Meningitis (G00,G03)	2	0.4	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	102	21.8	44.6
American Indian, female, 45-54 years				
...	All causes	738	100.0	342.4
1	Malignant neoplasms (C00-C97)	157	21.3	72.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	90	12.2	41.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	84	11.4	39.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	82	11.1	38.0
5	Diabetes mellitus (E10-E14)	47	6.4	21.8
6	Cerebrovascular diseases (I60-I69)	19	2.6	*
7	Septicemia (A40-A41)	18	2.4	*
7	Chronic lower respiratory diseases (J40-J47)	18	2.4	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	2.0	*
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	1.6	*
11	Viral hepatitis (B15-B19)	11	1.5	*
12	Influenza and pneumonia (J10-J18)	10	1.4	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	1.2	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.7	*
15	Pregnancy, childbirth and the puerperium (O00-099)	3	0.4	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.4	*
...	All other causes (Residual)	155	21.0	71.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 55-64 years				
...	All causes	932	100.0	686.6
1	Malignant neoplasms (C00-C97)	263	28.2	193.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	170	18.2	125.2
3	Diabetes mellitus (E10-E14)	87	9.3	64.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	53	5.7	39.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	43	4.6	31.7
6	Chronic lower respiratory diseases (J40-J47)	42	4.5	30.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	3.3	22.8
8	Cerebrovascular diseases (I60-I69)	22	2.4	16.2
9	Septicemia (A40-A41)	17	1.8	*
10	Influenza and pneumonia (J10-J18)	15	1.6	*
11	Viral hepatitis (B15-B19)	13	1.4	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	0.6	*
13	Pneumonitis due to solids and liquids (J69)	3	0.3	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.2	*
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.2	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	2	0.2	*
...	All other causes (Residual)	157	16.8	115.7
American Indian, female, 65-74 years				
...	All causes	1,157	100.0	1,657.3
1	Malignant neoplasms (C00-C97)	328	28.3	469.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	225	19.4	322.3
3	Diabetes mellitus (E10-E14)	108	9.3	154.7
4	Chronic lower respiratory diseases (J40-J47)	75	6.5	107.4
5	Cerebrovascular diseases (I60-I69)	55	4.8	78.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	3.9	64.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	36	3.1	51.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30	2.6	43.0
9	Septicemia (A40-A41)	16	1.4	*
10	Influenza and pneumonia (J10-J18)	13	1.1	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.5	*
12	Alzheimer's disease (G30)	6	0.5	*
12	Aortic aneurysm and dissection (I71)	6	0.5	*
12	Pneumonitis due to solids and liquids (J69)	6	0.5	*
...	All other causes (Residual)	193	16.7	276.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 75-84 years				
...	All causes	1,386	100.0	3,746.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	347	25.0	937.9
2	Malignant neoplasms (C00-C97)	280	20.2	756.8
3	Diabetes mellitus (E10-E14)	118	8.5	319.0
4	Cerebrovascular diseases (I60-I69)	99	7.1	267.6
5	Chronic lower respiratory diseases (J40-J47)	93	6.7	251.4
6	Alzheimer's disease (G30)	48	3.5	129.7
7	Influenza and pneumonia (J10-J18)	31	2.2	83.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29	2.1	78.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	2.0	75.7
10	Septicemia (A40-A41)	26	1.9	70.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	15	1.1	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	0.9	*
13	Pneumonitis due to solids and liquids (J69)	8	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.5	*
14	Parkinson's disease (G20-G21)	7	0.5	*
14	Atherosclerosis (I70)	7	0.5	*
...	All other causes (Residual)	231	16.7	624.4
American Indian, female, 85 years and over				
...	All causes	1,085	100.0	6,633.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	308	28.4	1,883.1
2	Malignant neoplasms (C00-C97)	112	10.3	684.8
3	Cerebrovascular diseases (I60-I69)	106	9.8	648.1
4	Diabetes mellitus (E10-E14)	60	5.5	366.8
5	Alzheimer's disease (G30)	52	4.8	317.9
6	Influenza and pneumonia (J10-J18)	42	3.9	256.8
7	Chronic lower respiratory diseases (J40-J47)	40	3.7	244.6
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	24	2.2	146.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	2.1	140.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	2.1	140.6
11	Septicemia (A40-A41)	12	1.1	*
11	Parkinson's disease (G20-G21)	12	1.1	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.0	*
14	Atherosclerosis (I70)	10	0.9	*
14	Pneumonitis due to solids and liquids (J69)	10	0.9	*
...	All other causes (Residual)	240	22.1	1,467.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, all ages				
...	All causes	44,707	100.0	307.4
1	Malignant neoplasms (C00-C97)	11,784	26.4	81.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,673	23.9	73.4
3	Cerebrovascular diseases (I60-I69)	3,662	8.2	25.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,125	4.8	14.6
5	Diabetes mellitus (E10-E14)	1,621	3.6	11.1
6	Chronic lower respiratory diseases (J40-J47)	1,352	3.0	9.3
7	Influenza and pneumonia (J10-J18)	1,347	3.0	9.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	866	1.9	6.0
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	813	1.8	5.6
10	Alzheimer's disease (G30)	720	1.6	5.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	585	1.3	4.0
12	Septicemia (A40-A41)	546	1.2	3.8
13	Certain conditions originating in the perinatal period (P00-P96)	454	1.0	3.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	427	1.0	2.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	419	0.9	2.9
...	All other causes (Residual)	7,313	16.4	50.3
Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	160	100.0	19.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	23.1	4.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	13.8	2.7
3	Malignant neoplasms (C00-C97)	19	11.9	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	6.9	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	4.4	*
6	Septicemia (A40-A41)	6	3.8	*
7	Influenza and pneumonia (J10-J18)	4	2.5	*
8	Pneumonitis due to solids and liquids (J69)	3	1.9	*
9	Chronic lower respiratory diseases (J40-J47)	2	1.3	*
9	Certain conditions originating in the perinatal period (P00-P96)	2	1.3	*
11	Tuberculosis (A16-A19)	1	0.6	*
11	Anemias (D50-D64)	1	0.6	*
11	Meningitis (G00,G03)	1	0.6	*
11	Cerebrovascular diseases (I60-I69)	1	0.6	*
11	Acute bronchitis and bronchiolitis (J20-J21)	1	0.6	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.6	*
...	All other causes (Residual)	41	25.6	5.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 5-14 years				
...	All causes	204	100.0	10.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	76	37.3	4.0
2	Malignant neoplasms (C00-C97)	37	18.1	2.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	6.4	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	6.4	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	5.9	*
6	Cerebrovascular diseases (I60-I69)	5	2.5	*
7	Influenza and pneumonia (J10-J18)	4	2.0	*
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3	1.5	*
9	Chronic lower respiratory diseases (J40-J47)	2	1.0	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	1.0	*
11	Septicemia (A40-A41)	1	0.5	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	*
11	Anemias (D50-D64)	1	0.5	*
11	Diseases of appendix (K35-K38)	1	0.5	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.5	*
...	All other causes (Residual)	32	15.7	1.7
Asian or Pacific Islander, both sexes, 15-24 years				
...	All causes	869	100.0	44.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	327	37.6	16.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	160	18.4	8.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	132	15.2	6.7
4	Malignant neoplasms (C00-C97)	70	8.1	3.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	2.8	1.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	2.1	*
7	Chronic lower respiratory diseases (J40-J47)	6	0.7	*
8	Septicemia (A40-A41)	5	0.6	*
9	Anemias (D50-D64)	4	0.5	*
9	Cerebrovascular diseases (I60-I69)	4	0.5	*
11	Diabetes mellitus (E10-E14)	3	0.3	*
11	Pregnancy, childbirth and the puerperium (O00-099)	3	0.3	*
11	Legal intervention (Y35,Y89.0)	3	0.3	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
15	Viral hepatitis (B15-B19)	2	0.2	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.2	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.2	*
15	Influenza and pneumonia (J10-J18)	2	0.2	*
...	All other causes (Residual)	99	11.4	5.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 25-34 years				
...	All causes	1,072	100.0	41.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	243	22.7	9.3
2	Malignant neoplasms (C00-C97)	186	17.4	7.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	175	16.3	6.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	96	9.0	3.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	86	8.0	3.3
6	Cerebrovascular diseases (I60-I69)	26	2.4	1.0
7	Pregnancy, childbirth and the puerperium (O00-099)	22	2.1	0.8
8	Influenza and pneumonia (J10-J18)	13	1.2	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	13	1.2	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	1.2	*
11	Diabetes mellitus (E10-E14)	12	1.1	*
12	Septicemia (A40-A41)	10	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.5	*
13	Anemias (D50-D64)	5	0.5	*
13	Aortic aneurysm and dissection (I71)	5	0.5	*
13	Chronic lower respiratory diseases (J40-J47)	5	0.5	*
...	All other causes (Residual)	157	14.6	6.0
Asian or Pacific Islander, both sexes, 35-44 years				
...	All causes	1,801	100.0	72.2
1	Malignant neoplasms (C00-C97)	554	30.8	22.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	241	13.4	9.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	240	13.3	9.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	142	7.9	5.7
5	Cerebrovascular diseases (I60-I69)	104	5.8	4.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	3.1	2.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	38	2.1	1.5
8	Diabetes mellitus (E10-E14)	31	1.7	1.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	27	1.5	1.1
10	Viral hepatitis (B15-B19)	22	1.2	0.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	1.2	0.8
12	Aortic aneurysm and dissection (I71)	15	0.8	*
12	Pregnancy, childbirth and the puerperium (O00-099)	15	0.8	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.7	*
15	Septicemia (A40-A41)	12	0.7	*
15	Chronic lower respiratory diseases (J40-J47)	12	0.7	*
...	All other causes (Residual)	258	14.3	10.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 45-54 years				
...	All causes	3,706	100.0	186.3
1	Malignant neoplasms (C00-C97)	1,431	38.6	71.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	689	18.6	34.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	261	7.0	13.1
4	Cerebrovascular diseases (I60-I69)	235	6.3	11.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	141	3.8	7.1
6	Diabetes mellitus (E10-E14)	112	3.0	5.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	97	2.6	4.9
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	71	1.9	3.6
9	Viral hepatitis (B15-B19)	59	1.6	3.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	1.0	1.9
11	Septicemia (A40-A41)	34	0.9	1.7
11	Influenza and pneumonia (J10-J18)	34	0.9	1.7
11	Chronic lower respiratory diseases (J40-J47)	34	0.9	1.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	31	0.8	1.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	26	0.7	1.3
...	All other causes (Residual)	414	11.2	20.8
Asian or Pacific Islander, both sexes, 55-64 years				
...	All causes	5,671	100.0	433.6
1	Malignant neoplasms (C00-C97)	2,319	40.9	177.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,220	21.5	93.3
3	Cerebrovascular diseases (I60-I69)	422	7.4	32.3
4	Diabetes mellitus (E10-E14)	260	4.6	19.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	218	3.8	16.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	95	1.7	7.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	87	1.5	6.7
8	Chronic lower respiratory diseases (J40-J47)	84	1.5	6.4
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	83	1.5	6.3
10	Viral hepatitis (B15-B19)	77	1.4	5.9
11	Septicemia (A40-A41)	66	1.2	5.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	64	1.1	4.9
13	Influenza and pneumonia (J10-J18)	60	1.1	4.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	0.7	2.8
15	Aortic aneurysm and dissection (I71)	27	0.5	2.1
...	All other causes (Residual)	552	9.7	42.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 65-74 years				
...	All causes	7,937	100.0	1,092.6
1	Malignant neoplasms (C00-C97)	2,827	35.6	389.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,876	23.6	258.2
3	Cerebrovascular diseases (I60-I69)	679	8.6	93.5
4	Diabetes mellitus (E10-E14)	370	4.7	50.9
5	Chronic lower respiratory diseases (J40-J47)	226	2.8	31.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	204	2.6	28.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	169	2.1	23.3
8	Influenza and pneumonia (J10-J18)	154	1.9	21.2
9	Septicemia (A40-A41)	114	1.4	15.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	85	1.1	11.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	80	1.0	11.0
12	Viral hepatitis (B15-B19)	69	0.9	9.5
13	Aortic aneurysm and dissection (I71)	63	0.8	8.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	59	0.7	8.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	58	0.7	8.0
...	All other causes (Residual)	904	11.4	124.4
Asian or Pacific Islander, both sexes, 75-84 years				
...	All causes	11,778	100.0	2,965.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,064	26.0	771.5
2	Malignant neoplasms (C00-C97)	2,942	25.0	740.8
3	Cerebrovascular diseases (I60-I69)	1,146	9.7	288.6
4	Chronic lower respiratory diseases (J40-J47)	547	4.6	137.7
5	Diabetes mellitus (E10-E14)	500	4.2	125.9
6	Influenza and pneumonia (J10-J18)	426	3.6	107.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	287	2.4	72.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	286	2.4	72.0
9	Alzheimer's disease (G30)	221	1.9	55.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	172	1.5	43.3
11	Septicemia (A40-A41)	167	1.4	42.1
12	Parkinson's disease (G20-G21)	130	1.1	32.7
13	Aortic aneurysm and dissection (I71)	111	0.9	27.9
14	Pneumonitis due to solids and liquids (J69)	88	0.7	22.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	87	0.7	21.9
...	All other causes (Residual)	1,604	13.6	403.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 85 years and over				
...	All causes	10,629	100.0	8,298.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,430	32.3	2,678.1
2	Malignant neoplasms (C00-C97)	1,395	13.1	1,089.2
3	Cerebrovascular diseases (I60-I69)	1,038	9.8	810.4
4	Influenza and pneumonia (J10-J18)	630	5.9	491.9
5	Alzheimer's disease (G30)	467	4.4	364.6
6	Chronic lower respiratory diseases (J40-J47)	431	4.1	336.5
7	Diabetes mellitus (E10-E14)	333	3.1	260.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	246	2.3	192.1
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	228	2.1	178.0
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	208	2.0	162.4
11	Septicemia (A40-A41)	123	1.2	96.0
12	Pneumonitis due to solids and liquids (J69)	120	1.1	93.7
13	Parkinson's disease (G20-G21)	119	1.1	92.9
14	Aortic aneurysm and dissection (I71)	75	0.7	58.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	67	0.6	52.3
...	All other causes (Residual)	1,719	16.2	1,342.2
Asian or Pacific Islander, male, all ages				
...	All causes	23,382	100.0	330.6
1	Malignant neoplasms (C00-C97)	5,974	25.5	84.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,827	24.9	82.4
3	Cerebrovascular diseases (I60-I69)	1,671	7.1	23.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,309	5.6	18.5
5	Chronic lower respiratory diseases (J40-J47)	847	3.6	12.0
6	Diabetes mellitus (E10-E14)	812	3.5	11.5
7	Influenza and pneumonia (J10-J18)	717	3.1	10.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	563	2.4	8.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	432	1.8	6.1
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	320	1.4	4.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	278	1.2	3.9
12	Certain conditions originating in the perinatal period (P00-P96)	257	1.1	3.6
13	Septicemia (A40-A41)	256	1.1	3.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	247	1.1	3.5
15	Alzheimer's disease (G30)	245	1.0	3.5
...	All other causes (Residual)	3,627	15.5	51.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 1-4 years				
...	All causes	76	100.0	18.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	30.3	5.5
2	Malignant neoplasms (C00-C97)	8	10.5	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	10.5	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	7.9	*
5	Septicemia (A40-A41)	2	2.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.6	*
5	Pneumonitis due to solids and liquids (J69)	2	2.6	*
8	Tuberculosis (A16-A19)	1	1.3	*
8	Anemias (D50-D64)	1	1.3	*
8	Meningitis (G00,G03)	1	1.3	*
8	Influenza and pneumonia (J10-J18)	1	1.3	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	1.3	*
...	All other causes (Residual)	20	26.3	4.8
Asian or Pacific Islander, male, 5-14 years				
...	All causes	108	100.0	11.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47	43.5	4.9
2	Malignant neoplasms (C00-C97)	20	18.5	2.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	8.3	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	4.6	*
5	Cerebrovascular diseases (I60-I69)	2	1.9	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.9	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	1.9	*
8	Septicemia (A40-A41)	1	0.9	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.9	*
8	Anemias (D50-D64)	1	0.9	*
8	Influenza and pneumonia (J10-J18)	1	0.9	*
8	Chronic lower respiratory diseases (J40-J47)	1	0.9	*
8	Diseases of appendix (K35-K38)	1	0.9	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.9	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.9	*
...	All other causes (Residual)	13	12.0	*

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 15-24 years				
...	All causes	624	100.0	61.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	243	38.9	24.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	121	19.4	12.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	116	18.6	11.5
4	Malignant neoplasms (C00-C97)	39	6.3	3.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	2.9	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	1.8	*
7	Chronic lower respiratory diseases (J40-J47)	5	0.8	*
8	Diabetes mellitus (E10-E14)	3	0.5	*
8	Cerebrovascular diseases (I60-I69)	3	0.5	*
8	Legal intervention (Y35,Y89.0)	3	0.5	*
11	Septicemia (A40-A41)	2	0.3	*
11	Viral hepatitis (B15-B19)	2	0.3	*
11	Anemias (D50-D64)	2	0.3	*
11	Influenza and pneumonia (J10-J18)	2	0.3	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	52	8.3	5.1
Asian or Pacific Islander, male, 25-34 years				
...	All causes	694	100.0	54.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	174	25.1	13.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	128	18.4	10.0
3	Malignant neoplasms (C00-C97)	89	12.8	6.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	75	10.8	5.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	9.2	5.0
6	Cerebrovascular diseases (I60-I69)	17	2.4	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	11	1.6	*
8	Diabetes mellitus (E10-E14)	9	1.3	*
9	Septicemia (A40-A41)	8	1.2	*
9	Influenza and pneumonia (J10-J18)	8	1.2	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	1.0	*
12	Chronic lower respiratory diseases (J40-J47)	5	0.7	*
13	Anemias (D50-D64)	3	0.4	*
13	Aortic aneurysm and dissection (I71)	3	0.4	*
13	Legal intervention (Y35,Y89.0)	3	0.4	*
...	All other causes (Residual)	90	13.0	7.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 35-44 years				
...	All causes	1,075	100.0	88.5
1	Malignant neoplasms (C00-C97)	238	22.1	19.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	192	17.9	15.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	159	14.8	13.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	102	9.5	8.4
5	Cerebrovascular diseases (I60-I69)	51	4.7	4.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	3.3	3.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	28	2.6	2.3
8	Human immunodeficiency virus (HIV) disease (B20-B24)	23	2.1	1.9
9	Viral hepatitis (B15-B19)	20	1.9	1.6
9	Diabetes mellitus (E10-E14)	20	1.9	1.6
11	Aortic aneurysm and dissection (I71)	13	1.2	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	1.2	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.7	*
14	Septicemia (A40-A41)	7	0.7	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.6	*
...	All other causes (Residual)	159	14.8	13.1
Asian or Pacific Islander, male, 45-54 years				
...	All causes	2,180	100.0	232.5
1	Malignant neoplasms (C00-C97)	658	30.2	70.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	522	23.9	55.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	173	7.9	18.4
4	Cerebrovascular diseases (I60-I69)	126	5.8	13.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	94	4.3	10.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	84	3.9	9.0
7	Diabetes mellitus (E10-E14)	72	3.3	7.7
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	47	2.2	5.0
9	Viral hepatitis (B15-B19)	43	2.0	4.6
10	Human immunodeficiency virus (HIV) disease (B20-B24)	26	1.2	2.8
11	Influenza and pneumonia (J10-J18)	22	1.0	2.3
12	Chronic lower respiratory diseases (J40-J47)	20	0.9	2.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	0.9	2.1
14	Septicemia (A40-A41)	18	0.8	*
15	Aortic aneurysm and dissection (I71)	15	0.7	*
...	All other causes (Residual)	240	11.0	25.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 55-64 years				
...	All causes	3,334	100.0	550.7
1	Malignant neoplasms (C00-C97)	1,194	35.8	197.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	880	26.4	145.4
3	Cerebrovascular diseases (I60-I69)	220	6.6	36.3
4	Diabetes mellitus (E10-E14)	167	5.0	27.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	129	3.9	21.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	68	2.0	11.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	56	1.7	9.3
8	Chronic lower respiratory diseases (J40-J47)	53	1.6	8.8
9	Viral hepatitis (B15-B19)	50	1.5	8.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	1.3	7.4
11	Septicemia (A40-A41)	38	1.1	6.3
11	Influenza and pneumonia (J10-J18)	38	1.1	6.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	35	1.0	5.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	26	0.8	4.3
15	Aortic aneurysm and dissection (I71)	20	0.6	3.3
...	All other causes (Residual)	315	9.4	52.0
Asian or Pacific Islander, male, 65-74 years				
...	All causes	4,358	100.0	1,329.2
1	Malignant neoplasms (C00-C97)	1,508	34.6	459.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,129	25.9	344.3
3	Cerebrovascular diseases (I60-I69)	357	8.2	108.9
4	Diabetes mellitus (E10-E14)	189	4.3	57.6
5	Chronic lower respiratory diseases (J40-J47)	146	3.4	44.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	107	2.5	32.6
7	Influenza and pneumonia (J10-J18)	96	2.2	29.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	92	2.1	28.1
9	Septicemia (A40-A41)	55	1.3	16.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	45	1.0	13.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	43	1.0	13.1
12	Aortic aneurysm and dissection (I71)	40	0.9	12.2
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	0.8	10.1
14	Viral hepatitis (B15-B19)	31	0.7	9.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.7	8.8
15	Parkinson's disease (G20-G21)	29	0.7	8.8
...	All other causes (Residual)	429	9.8	130.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 75-84 years				
...	All causes	5,833	100.0	3,606.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,558	26.7	963.3
2	Malignant neoplasms (C00-C97)	1,524	26.1	942.3
3	Cerebrovascular diseases (I60-I69)	477	8.2	294.9
4	Chronic lower respiratory diseases (J40-J47)	355	6.1	219.5
5	Influenza and pneumonia (J10-J18)	228	3.9	141.0
6	Diabetes mellitus (E10-E14)	224	3.8	138.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	143	2.5	88.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	138	2.4	85.3
9	Parkinson's disease (G20-G21)	79	1.4	48.8
10	Alzheimer's disease (G30)	78	1.3	48.2
11	Septicemia (A40-A41)	73	1.3	45.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	72	1.2	44.5
13	Aortic aneurysm and dissection (I71)	62	1.1	38.3
14	Pneumonitis due to solids and liquids (J69)	48	0.8	29.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	0.6	22.3
...	All other causes (Residual)	738	12.7	456.3
Asian or Pacific Islander, male, 85 years and over				
...	All causes	4,587	100.0	9,524.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,438	31.3	2,985.9
2	Malignant neoplasms (C00-C97)	693	15.1	1,439.0
3	Cerebrovascular diseases (I60-I69)	417	9.1	865.9
4	Influenza and pneumonia (J10-J18)	311	6.8	645.8
5	Chronic lower respiratory diseases (J40-J47)	255	5.6	529.5
6	Alzheimer's disease (G30)	155	3.4	321.9
7	Diabetes mellitus (E10-E14)	128	2.8	265.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	118	2.6	245.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	98	2.1	203.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	75	1.6	155.7
11	Pneumonitis due to solids and liquids (J69)	65	1.4	135.0
12	Parkinson's disease (G20-G21)	64	1.4	132.9
13	Septicemia (A40-A41)	48	1.0	99.7
14	Aortic aneurysm and dissection (I71)	45	1.0	93.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	0.8	74.8
...	All other causes (Residual)	641	14.0	1,331.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, all ages				
...	All causes	21,325	100.0	285.6
1	Malignant neoplasms (C00-C97)	5,810	27.2	77.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,846	22.7	64.9
3	Cerebrovascular diseases (I60-I69)	1,991	9.3	26.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	816	3.8	10.9
5	Diabetes mellitus (E10-E14)	809	3.8	10.8
6	Influenza and pneumonia (J10-J18)	630	3.0	8.4
7	Chronic lower respiratory diseases (J40-J47)	505	2.4	6.8
8	Alzheimer's disease (G30)	475	2.2	6.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	434	2.0	5.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	338	1.6	4.5
11	Septicemia (A40-A41)	290	1.4	3.9
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	250	1.2	3.3
13	Certain conditions originating in the perinatal period (P00-P96)	197	0.9	2.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	146	0.7	2.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	141	0.7	1.9
...	All other causes (Residual)	3,647	17.1	48.8
Asian or Pacific Islander, female, 1-4 years				
...	All causes	84	100.0	21.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	16.7	*
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	16.7	*
3	Malignant neoplasms (C00-C97)	11	13.1	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	6.0	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	6.0	*
6	Septicemia (A40-A41)	4	4.8	*
7	Influenza and pneumonia (J10-J18)	3	3.6	*
8	Chronic lower respiratory diseases (J40-J47)	2	2.4	*
9	Cerebrovascular diseases (I60-I69)	1	1.2	*
9	Acute bronchitis and bronchiolitis (J20-J21)	1	1.2	*
9	Pneumonitis due to solids and liquids (J69)	1	1.2	*
9	Certain conditions originating in the perinatal period (P00-P96)	1	1.2	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.2	*
...	All other causes (Residual)	21	25.0	5.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 5-14 years				
...	All causes	96	100.0	10.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29	30.2	3.1
2	Malignant neoplasms (C00-C97)	17	17.7	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	11.5	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	8.3	*
5	Cerebrovascular diseases (I60-I69)	3	3.1	*
5	Influenza and pneumonia (J10-J18)	3	3.1	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	3.1	*
8	Chronic lower respiratory diseases (J40-J47)	1	1.0	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.0	*
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	1.0	*
...	All other causes (Residual)	19	19.8	*
Asian or Pacific Islander, female, 15-24 years				
...	All causes	245	100.0	25.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	84	34.3	8.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	15.9	4.0
3	Malignant neoplasms (C00-C97)	31	12.7	3.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	6.5	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	2.9	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	2.4	*
7	Septicemia (A40-A41)	3	1.2	*
7	Pregnancy, childbirth and the puerperium (O00-099)	3	1.2	*
9	Anemias (D50-D64)	2	0.8	*
10	Syphilis (A50-A53)	1	0.4	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	*
10	Cerebrovascular diseases (I60-I69)	1	0.4	*
10	Chronic lower respiratory diseases (J40-J47)	1	0.4	*
10	Hernia (K40-K46)	1	0.4	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
10	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.4	*
...	All other causes (Residual)	46	18.8	4.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 25-34 years				
...	All causes	378	100.0	28.5
1	Malignant neoplasms (C00-C97)	97	25.7	7.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	69	18.3	5.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47	12.4	3.5
4	Pregnancy, childbirth and the puerperium (O00-099)	22	5.8	1.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	5.8	1.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	5.6	1.6
7	Cerebrovascular diseases (I60-I69)	9	2.4	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	1.6	*
9	Influenza and pneumonia (J10-J18)	5	1.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.8	*
10	Diabetes mellitus (E10-E14)	3	0.8	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.8	*
13	Septicemia (A40-A41)	2	0.5	*
13	Anemias (D50-D64)	2	0.5	*
13	Aortic aneurysm and dissection (I71)	2	0.5	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.5	*
...	All other causes (Residual)	63	16.7	4.7
Asian or Pacific Islander, female, 35-44 years				
...	All causes	726	100.0	56.8
1	Malignant neoplasms (C00-C97)	316	43.5	24.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	81	11.2	6.3
3	Cerebrovascular diseases (I60-I69)	53	7.3	4.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	6.7	3.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	5.5	3.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	2.8	1.6
7	Pregnancy, childbirth and the puerperium (O00-099)	15	2.1	*
8	Diabetes mellitus (E10-E14)	11	1.5	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	10	1.4	*
10	Chronic lower respiratory diseases (J40-J47)	8	1.1	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	1.1	*
12	Influenza and pneumonia (J10-J18)	6	0.8	*
13	Septicemia (A40-A41)	5	0.7	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.7	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.6	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	0.6	*
...	All other causes (Residual)	91	12.5	7.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 45-54 years				
...	All causes	1,526	100.0	145.2
1	Malignant neoplasms (C00-C97)	773	50.7	73.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	167	10.9	15.9
3	Cerebrovascular diseases (I60-I69)	109	7.1	10.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	88	5.8	8.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47	3.1	4.5
6	Diabetes mellitus (E10-E14)	40	2.6	3.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	1.6	2.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	1.1	*
9	Septicemia (A40-A41)	16	1.0	*
9	Viral hepatitis (B15-B19)	16	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	14	0.9	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	13	0.9	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	0.8	*
13	Influenza and pneumonia (J10-J18)	12	0.8	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.7	*
...	All other causes (Residual)	167	10.9	15.9
Asian or Pacific Islander, female, 55-64 years				
...	All causes	2,337	100.0	332.7
1	Malignant neoplasms (C00-C97)	1,125	48.1	160.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	340	14.5	48.4
3	Cerebrovascular diseases (I60-I69)	202	8.6	28.8
4	Diabetes mellitus (E10-E14)	93	4.0	13.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	89	3.8	12.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	2.1	7.1
7	Chronic lower respiratory diseases (J40-J47)	31	1.3	4.4
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	29	1.2	4.1
9	Septicemia (A40-A41)	28	1.2	4.0
10	Viral hepatitis (B15-B19)	27	1.2	3.8
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27	1.2	3.8
12	Influenza and pneumonia (J10-J18)	22	0.9	3.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	19	0.8	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.7	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	0.5	*
...	All other causes (Residual)	227	9.7	32.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 65-74 years				
...	All causes	3,579	100.0	897.9
1	Malignant neoplasms (C00-C97)	1,319	36.9	330.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	747	20.9	187.4
3	Cerebrovascular diseases (I60-I69)	322	9.0	80.8
4	Diabetes mellitus (E10-E14)	181	5.1	45.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	97	2.7	24.3
6	Chronic lower respiratory diseases (J40-J47)	80	2.2	20.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	77	2.2	19.3
8	Septicemia (A40-A41)	59	1.6	14.8
9	Influenza and pneumonia (J10-J18)	58	1.6	14.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	40	1.1	10.0
11	Viral hepatitis (B15-B19)	38	1.1	9.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	37	1.0	9.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.8	7.5
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25	0.7	6.3
15	Aortic aneurysm and dissection (I71)	23	0.6	5.8
...	All other causes (Residual)	446	12.5	111.9
Asian or Pacific Islander, female, 75-84 years				
...	All causes	5,945	100.0	2,525.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,506	25.3	639.8
2	Malignant neoplasms (C00-C97)	1,418	23.9	602.4
3	Cerebrovascular diseases (I60-I69)	669	11.3	284.2
4	Diabetes mellitus (E10-E14)	276	4.6	117.2
5	Influenza and pneumonia (J10-J18)	198	3.3	84.1
6	Chronic lower respiratory diseases (J40-J47)	192	3.2	81.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	148	2.5	62.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	144	2.4	61.2
9	Alzheimer's disease (G30)	143	2.4	60.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	100	1.7	42.5
11	Septicemia (A40-A41)	94	1.6	39.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	52	0.9	22.1
13	Parkinson's disease (G20-G21)	51	0.9	21.7
14	Aortic aneurysm and dissection (I71)	49	0.8	20.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	42	0.7	17.8
...	All other causes (Residual)	863	14.5	366.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
	Asian or Pacific Islander, female, 85 years and over			
...	All causes	6,042	100.0	7,560.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,992	33.0	2,492.6
2	Malignant neoplasms (C00-C97)	702	11.6	878.4
3	Cerebrovascular diseases (I60-I69)	621	10.3	777.0
4	Influenza and pneumonia (J10-J18)	319	5.3	399.2
5	Alzheimer's disease (G30)	312	5.2	390.4
6	Diabetes mellitus (E10-E14)	205	3.4	256.5
7	Chronic lower respiratory diseases (J40-J47)	176	2.9	220.2
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	153	2.5	191.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	128	2.1	160.2
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	110	1.8	137.6
11	Septicemia (A40-A41)	75	1.2	93.8
12	Parkinson's disease (G20-G21)	55	0.9	68.8
12	Pneumonitis due to solids and liquids (J69)	55	0.9	68.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	0.5	38.8
15	Aortic aneurysm and dissection (I71)	30	0.5	37.5
...	All other causes (Residual)	1,078	17.8	1,348.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable