

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2011

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Hispanic, both sexes				
...	All causes	4,823	100.0	525.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,282	26.6	139.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	782	16.2	85.2
3	Newborn affected by maternal complications of pregnancy (P01)	329	6.8	35.8
4	Sudden infant death syndrome (R95)	261	5.4	28.4
5	Newborn affected by complications of placenta, cord and membranes (P02)	202	4.2	22.0
6	Accidents (unintentional injuries) (V01-X59)	159	3.3	17.3
7	Bacterial sepsis of newborn (P36)	109	2.3	11.9
8	Diseases of the circulatory system (I00-I99)	107	2.2	11.7
9	Respiratory distress of newborn (P22)	101	2.1	11.0
10	Neonatal hemorrhage (P50-P52,P54)	95	2.0	10.3
11	Necrotizing enterocolitis of newborn (P77)	76	1.6	8.3
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	66	1.4	7.2
13	Atelectasis (P28.0-P28.1)	59	1.2	6.4
14	Assault (homicide) (*U01,X85-Y09)	51	1.1	5.6
15	Diarrhea and gastroenteritis of infectious origin (A09)	39	0.8	4.2
15	Hydrops fetalis not due to hemolytic disease (P83.2)	39	0.8	4.2
...	All other causes (Residual)	1,066	22.1	116.1
Infants, Hispanic, male				
...	All causes	2,618	100.0	559.2
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	665	25.4	142.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	422	16.1	90.1
3	Newborn affected by maternal complications of pregnancy (P01)	178	6.8	38.0
4	Sudden infant death syndrome (R95)	148	5.7	31.6
5	Newborn affected by complications of placenta, cord and membranes (P02)	120	4.6	25.6
6	Accidents (unintentional injuries) (V01-X59)	89	3.4	19.0
7	Respiratory distress of newborn (P22)	58	2.2	12.4
8	Diseases of the circulatory system (I00-I99)	56	2.1	12.0
8	Bacterial sepsis of newborn (P36)	56	2.1	12.0
10	Neonatal hemorrhage (P50-P52,P54)	54	2.1	11.5
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	41	1.6	8.8
12	Necrotizing enterocolitis of newborn (P77)	39	1.5	8.3
13	Atelectasis (P28.0-P28.1)	33	1.3	7.0
14	Assault (homicide) (*U01,X85-Y09)	31	1.2	6.6
15	Diarrhea and gastroenteritis of infectious origin (A09)	24	0.9	5.1
...	All other causes (Residual)	604	23.1	129.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2011

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Hispanic, female				
...	All causes	2,205	100.0	490.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	617	28.0	137.1
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	360	16.3	80.0
3	Newborn affected by maternal complications of pregnancy (P01)	151	6.8	33.6
4	Sudden infant death syndrome (R95)	113	5.1	25.1
5	Newborn affected by complications of placenta, cord and membranes (P02)	82	3.7	18.2
6	Accidents (unintentional injuries) (V01-X59)	70	3.2	15.6
7	Bacterial sepsis of newborn (P36)	53	2.4	11.8
8	Diseases of the circulatory system (I00-I99)	51	2.3	11.3
9	Respiratory distress of newborn (P22)	43	2.0	9.6
10	Neonatal hemorrhage (P50-P52,P54)	41	1.9	9.1
11	Necrotizing enterocolitis of newborn (P77)	37	1.7	8.2
12	Atelectasis (P28.0-P28.1)	26	1.2	5.8
13	Intrauterine hypoxia and birth asphyxia (P20-P21)	25	1.1	5.6
14	Assault (homicide) (*U01,X85-Y09)	20	0.9	4.4
15	Hydrops fetalis not due to hemolytic disease (P83.2)	19	0.9	*
...	All other causes (Residual)	497	22.5	110.4
Infants, Non-Hispanic, white, both sexes				
...	All causes	10,842	100.0	505.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,515	23.2	117.2
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,548	14.3	72.1
3	Sudden infant death syndrome (R95)	1,000	9.2	46.6
4	Newborn affected by maternal complications of pregnancy (P01)	640	5.9	29.8
5	Accidents (unintentional injuries) (V01-X59)	579	5.3	27.0
6	Newborn affected by complications of placenta, cord and membranes (P02)	456	4.2	21.2
7	Respiratory distress of newborn (P22)	239	2.2	11.1
8	Neonatal hemorrhage (P50-P52,P54)	233	2.1	10.9
9	Bacterial sepsis of newborn (P36)	222	2.0	10.3
10	Diseases of the circulatory system (I00-I99)	214	2.0	10.0
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	161	1.5	7.5
12	Necrotizing enterocolitis of newborn (P77)	145	1.3	6.8
13	Atelectasis (P28.0-P28.1)	126	1.2	5.9
14	Assault (homicide) (*U01,X85-Y09)	124	1.1	5.8
15	Hydrops fetalis not due to hemolytic disease (P83.2)	102	0.9	4.8
...	All other causes (Residual)	2,538	23.4	118.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2011

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Non-Hispanic, white, male				
...	All causes	6,082	100.0	551.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,328	21.8	120.5
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	859	14.1	77.9
3	Sudden infant death syndrome (R95)	586	9.6	53.2
4	Newborn affected by maternal complications of pregnancy (P01)	380	6.2	34.5
5	Accidents (unintentional injuries) (V01-X59)	323	5.3	29.3
6	Newborn affected by complications of placenta, cord and membranes (P02)	233	3.8	21.1
7	Neonatal hemorrhage (P50-P52,P54)	137	2.3	12.4
8	Respiratory distress of newborn (P22)	134	2.2	12.2
9	Bacterial sepsis of newborn (P36)	128	2.1	11.6
10	Diseases of the circulatory system (I00-I99)	119	2.0	10.8
11	Necrotizing enterocolitis of newborn (P77)	83	1.4	7.5
12	Atelectasis (P28.0-P28.1)	82	1.3	7.4
13	Intrauterine hypoxia and birth asphyxia (P20-P21)	75	1.2	6.8
14	Assault (homicide) (*U01,X85-Y09)	73	1.2	6.6
15	Influenza and pneumonia (J09-J18)	55	0.9	5.0
...	All other causes (Residual)	1,487	24.4	134.9
Infants, Non-Hispanic, white, female				
...	All causes	4,760	100.0	455.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,187	24.9	113.7
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	689	14.5	66.0
3	Sudden infant death syndrome (R95)	414	8.7	39.6
4	Newborn affected by maternal complications of pregnancy (P01)	260	5.5	24.9
5	Accidents (unintentional injuries) (V01-X59)	256	5.4	24.5
6	Newborn affected by complications of placenta, cord and membranes (P02)	223	4.7	21.4
7	Respiratory distress of newborn (P22)	105	2.2	10.1
8	Neonatal hemorrhage (P50-P52,P54)	96	2.0	9.2
9	Diseases of the circulatory system (I00-I99)	95	2.0	9.1
10	Bacterial sepsis of newborn (P36)	94	2.0	9.0
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	86	1.8	8.2
12	Necrotizing enterocolitis of newborn (P77)	62	1.3	5.9
13	Hydrops fetalis not due to hemolytic disease (P83.2)	53	1.1	5.1
14	Assault (homicide) (*U01,X85-Y09)	51	1.1	4.9
15	Atelectasis (P28.0-P28.1)	44	0.9	4.2
...	All other causes (Residual)	1,045	22.0	100.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2011

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Non-Hispanic, black, both sexes				
...	All causes	6,979	100.0	1,198.4
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,521	21.8	261.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	959	13.7	164.7
3	Sudden infant death syndrome (R95)	566	8.1	97.2
4	Newborn affected by maternal complications of pregnancy (P01)	522	7.5	89.6
5	Accidents (unintentional injuries) (V01-X59)	364	5.2	62.5
6	Newborn affected by complications of placenta, cord and membranes (P02)	307	4.4	52.7
7	Bacterial sepsis of newborn (P36)	172	2.5	29.5
8	Diseases of the circulatory system (I00-I99)	150	2.1	25.8
9	Necrotizing enterocolitis of newborn (P77)	149	2.1	25.6
10	Respiratory distress of newborn (P22)	139	2.0	23.9
11	Diarrhea and gastroenteritis of infectious origin (A09)	118	1.7	20.3
12	Neonatal hemorrhage (P50-P52,P54)	110	1.6	18.9
13	Assault (homicide) (*U01,X85-Y09)	94	1.3	16.1
14	Influenza and pneumonia (J09-J18)	80	1.1	13.7
15	Pulmonary hemorrhage originating in the perinatal period (P26)	77	1.1	13.2
...	All other causes (Residual)	1,651	23.7	283.5
Infants, Non-Hispanic, black, male				
...	All causes	3,885	100.0	1,312.7
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	858	22.1	289.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	504	13.0	170.3
3	Sudden infant death syndrome (R95)	321	8.3	108.5
4	Newborn affected by maternal complications of pregnancy (P01)	296	7.6	100.0
5	Accidents (unintentional injuries) (V01-X59)	199	5.1	67.2
6	Newborn affected by complications of placenta, cord and membranes (P02)	169	4.4	57.1
7	Bacterial sepsis of newborn (P36)	95	2.4	32.1
8	Necrotizing enterocolitis of newborn (P77)	84	2.2	28.4
9	Diseases of the circulatory system (I00-I99)	78	2.0	26.4
10	Diarrhea and gastroenteritis of infectious origin (A09)	71	1.8	24.0
11	Respiratory distress of newborn (P22)	67	1.7	22.6
12	Neonatal hemorrhage (P50-P52,P54)	62	1.6	20.9
13	Assault (homicide) (*U01,X85-Y09)	56	1.4	18.9
14	Pulmonary hemorrhage originating in the perinatal period (P26)	49	1.3	16.6
15	Influenza and pneumonia (J09-J18)	45	1.2	15.2
...	All other causes (Residual)	931	24.0	314.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2011

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Non-Hispanic, black, female				
...	All causes	3,094	100.0	1,080.3
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	663	21.4	231.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	455	14.7	158.9
3	Sudden infant death syndrome (R95)	245	7.9	85.5
4	Newborn affected by maternal complications of pregnancy (P01)	226	7.3	78.9
5	Accidents (unintentional injuries) (V01-X59)	165	5.3	57.6
6	Newborn affected by complications of placenta, cord and membranes (P02)	138	4.5	48.2
7	Bacterial sepsis of newborn (P36)	77	2.5	26.9
8	Diseases of the circulatory system (I00-I99)	72	2.3	25.1
8	Respiratory distress of newborn (P22)	72	2.3	25.1
10	Necrotizing enterocolitis of newborn (P77)	65	2.1	22.7
11	Neonatal hemorrhage (P50-P52,P54)	48	1.6	16.8
12	Diarrhea and gastroenteritis of infectious origin (A09)	47	1.5	16.4
13	Intrauterine hypoxia and birth asphyxia (P20-P21)	42	1.4	14.7
14	Assault (homicide) (*U01,X85-Y09)	38	1.2	13.3
15	Influenza and pneumonia (J09-J18)	35	1.1	12.2
...	All other causes (Residual)	706	22.8	246.5
Neonates, Hispanic, both sexes				
...	All causes	3,372	100.0	367.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	920	27.3	100.2
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	764	22.7	83.2
3	Newborn affected by maternal complications of pregnancy (P01)	326	9.7	35.5
4	Newborn affected by complications of placenta, cord and membranes (P02)	201	6.0	21.9
5	Bacterial sepsis of newborn (P36)	103	3.1	11.2
6	Respiratory distress of newborn (P22)	98	2.9	10.7
7	Neonatal hemorrhage (P50-P52,P54)	91	2.7	9.9
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	64	1.9	7.0
9	Necrotizing enterocolitis of newborn (P77)	62	1.8	6.8
10	Atelectasis (P28.0-P28.1)	59	1.7	6.4
11	Hydrops fetalis not due to hemolytic disease (P83.2)	38	1.1	4.1
12	Pulmonary hemorrhage originating in the perinatal period (P26)	31	0.9	3.4
13	Sudden infant death syndrome (R95)	24	0.7	2.6
14	Diseases of the circulatory system (I00-I99)	22	0.7	2.4
15	Newborn affected by maternal hypertensive disorders (P00.0)	21	0.6	2.3
...	All other causes (Residual)	548	16.3	59.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2011

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Hispanic, male				
...	All causes	1,813	100.0	387.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	489	27.0	104.5
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	412	22.7	88.0
3	Newborn affected by maternal complications of pregnancy (P01)	176	9.7	37.6
4	Newborn affected by complications of placenta, cord and membranes (P02)	120	6.6	25.6
5	Respiratory distress of newborn (P22)	55	3.0	11.7
6	Bacterial sepsis of newborn (P36)	53	2.9	11.3
7	Neonatal hemorrhage (P50-P52,P54)	50	2.8	10.7
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	40	2.2	8.5
9	Atelectasis (P28.0-P28.1)	33	1.8	7.0
10	Necrotizing enterocolitis of newborn (P77)	32	1.8	6.8
11	Hydrops fetalis not due to hemolytic disease (P83.2)	19	1.0	*
12	Pulmonary hemorrhage originating in the perinatal period (P26)	17	0.9	*
13	Congenital pneumonia (P23)	11	0.6	*
13	Sudden infant death syndrome (R95)	11	0.6	*
15	Newborn affected by other complications of labor and delivery (P03)	10	0.6	*
...	All other causes (Residual)	285	15.7	60.9
Neonates, Hispanic, female				
...	All causes	1,559	100.0	346.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	431	27.6	95.8
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	352	22.6	78.2
3	Newborn affected by maternal complications of pregnancy (P01)	150	9.6	33.3
4	Newborn affected by complications of placenta, cord and membranes (P02)	81	5.2	18.0
5	Bacterial sepsis of newborn (P36)	50	3.2	11.1
6	Respiratory distress of newborn (P22)	43	2.8	9.6
7	Neonatal hemorrhage (P50-P52,P54)	41	2.6	9.1
8	Necrotizing enterocolitis of newborn (P77)	30	1.9	6.7
9	Atelectasis (P28.0-P28.1)	26	1.7	5.8
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	24	1.5	5.3
11	Hydrops fetalis not due to hemolytic disease (P83.2)	19	1.2	*
12	Diseases of the circulatory system (I00-I99)	14	0.9	*
12	Pulmonary hemorrhage originating in the perinatal period (P26)	14	0.9	*
14	Sudden infant death syndrome (R95)	13	0.8	*
15	Newborn affected by maternal hypertensive disorders (P00.0)	12	0.8	*
...	All other causes (Residual)	259	16.6	57.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2011

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Non-Hispanic, white, both sexes				
...	All causes	7,176	100.0	334.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,814	25.3	84.5
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,509	21.0	70.3
3	Newborn affected by maternal complications of pregnancy (P01)	635	8.8	29.6
4	Newborn affected by complications of placenta, cord and membranes (P02)	450	6.3	21.0
5	Respiratory distress of newborn (P22)	238	3.3	11.1
6	Neonatal hemorrhage (P50-P52,P54)	229	3.2	10.7
7	Bacterial sepsis of newborn (P36)	211	2.9	9.8
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	158	2.2	7.4
9	Necrotizing enterocolitis of newborn (P77)	131	1.8	6.1
10	Atelectasis (P28.0-P28.1)	117	1.6	5.5
11	Sudden infant death syndrome (R95)	101	1.4	4.7
12	Hydrops fetalis not due to hemolytic disease (P83.2)	96	1.3	4.5
13	Accidents (unintentional injuries) (V01-X59)	78	1.1	3.6
14	Newborn affected by other complications of labor and delivery (P03)	71	1.0	3.3
15	Diseases of the circulatory system (I00-I99)	64	0.9	3.0
...	All other causes (Residual)	1,274	17.8	59.4
Neonates, Non-Hispanic, white, male				
...	All causes	3,985	100.0	361.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	973	24.4	88.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	838	21.0	76.0
3	Newborn affected by maternal complications of pregnancy (P01)	379	9.5	34.4
4	Newborn affected by complications of placenta, cord and membranes (P02)	230	5.8	20.9
5	Respiratory distress of newborn (P22)	134	3.4	12.2
5	Neonatal hemorrhage (P50-P52,P54)	134	3.4	12.2
7	Bacterial sepsis of newborn (P36)	121	3.0	11.0
8	Atelectasis (P28.0-P28.1)	77	1.9	7.0
8	Necrotizing enterocolitis of newborn (P77)	77	1.9	7.0
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	73	1.8	6.6
11	Sudden infant death syndrome (R95)	57	1.4	5.2
12	Hydrops fetalis not due to hemolytic disease (P83.2)	46	1.2	4.2
13	Newborn affected by other complications of labor and delivery (P03)	43	1.1	3.9
14	Accidents (unintentional injuries) (V01-X59)	38	1.0	3.4
15	Diseases of the circulatory system (I00-I99)	35	0.9	3.2
...	All other causes (Residual)	730	18.3	66.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2011

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Non-Hispanic, white, female				
...	All causes	3,191	100.0	305.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	841	26.4	80.5
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	671	21.0	64.2
3	Newborn affected by maternal complications of pregnancy (P01)	256	8.0	24.5
4	Newborn affected by complications of placenta, cord and membranes (P02)	220	6.9	21.1
5	Respiratory distress of newborn (P22)	104	3.3	10.0
6	Neonatal hemorrhage (P50-P52,P54)	95	3.0	9.1
7	Bacterial sepsis of newborn (P36)	90	2.8	8.6
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	85	2.7	8.1
9	Necrotizing enterocolitis of newborn (P77)	54	1.7	5.2
10	Hydrops fetalis not due to hemolytic disease (P83.2)	50	1.6	4.8
11	Sudden infant death syndrome (R95)	44	1.4	4.2
12	Atelectasis (P28.0-P28.1)	40	1.3	3.8
12	Accidents (unintentional injuries) (V01-X59)	40	1.3	3.8
14	Diseases of the circulatory system (I00-I99)	29	0.9	2.8
15	Newborn affected by other complications of labor and delivery (P03)	28	0.9	2.7
...	All other causes (Residual)	544	17.0	52.1
Neonates, Non-Hispanic, black, both sexes				
...	All causes	4,569	100.0	784.6
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,474	32.3	253.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	639	14.0	109.7
3	Newborn affected by maternal complications of pregnancy (P01)	520	11.4	89.3
4	Newborn affected by complications of placenta, cord and membranes (P02)	300	6.6	51.5
5	Bacterial sepsis of newborn (P36)	165	3.6	28.3
6	Respiratory distress of newborn (P22)	135	3.0	23.2
7	Necrotizing enterocolitis of newborn (P77)	132	2.9	22.7
8	Neonatal hemorrhage (P50-P52,P54)	109	2.4	18.7
9	Pulmonary hemorrhage originating in the perinatal period (P26)	76	1.7	13.1
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	70	1.5	12.0
11	Atelectasis (P28.0-P28.1)	64	1.4	11.0
12	Sudden infant death syndrome (R95)	42	0.9	7.2
13	Newborn affected by other complications of labor and delivery (P03)	31	0.7	5.3
14	Interstitial emphysema and related conditions originating in the perinatal period (P25)	28	0.6	4.8
15	Accidents (unintentional injuries) (V01-X59)	27	0.6	4.6
...	All other causes (Residual)	757	16.6	130.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2011

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Non-Hispanic, black, male				
...	All causes	2,525	100.0	853.2
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	827	32.8	279.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	339	13.4	114.5
3	Newborn affected by maternal complications of pregnancy (P01)	296	11.7	100.0
4	Newborn affected by complications of placenta, cord and membranes (P02)	167	6.6	56.4
5	Bacterial sepsis of newborn (P36)	91	3.6	30.7
6	Necrotizing enterocolitis of newborn (P77)	77	3.0	26.0
7	Respiratory distress of newborn (P22)	66	2.6	22.3
8	Neonatal hemorrhage (P50-P52,P54)	61	2.4	20.6
9	Pulmonary hemorrhage originating in the perinatal period (P26)	48	1.9	16.2
10	Atelectasis (P28.0-P28.1)	34	1.3	11.5
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	32	1.3	10.8
12	Sudden infant death syndrome (R95)	18	0.7	*
13	Accidents (unintentional injuries) (V01-X59)	16	0.6	*
14	Interstitial emphysema and related conditions originating in the perinatal period (P25)	15	0.6	*
15	Newborn affected by other complications of labor and delivery (P03)	14	0.6	*
...	All other causes (Residual)	424	16.8	143.3
Neonates, Non-Hispanic, black, female				
...	All causes	2,044	100.0	713.7
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	647	31.7	225.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	300	14.7	104.7
3	Newborn affected by maternal complications of pregnancy (P01)	224	11.0	78.2
4	Newborn affected by complications of placenta, cord and membranes (P02)	133	6.5	46.4
5	Bacterial sepsis of newborn (P36)	74	3.6	25.8
6	Respiratory distress of newborn (P22)	69	3.4	24.1
7	Necrotizing enterocolitis of newborn (P77)	55	2.7	19.2
8	Neonatal hemorrhage (P50-P52,P54)	48	2.3	16.8
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	38	1.9	13.3
10	Atelectasis (P28.0-P28.1)	30	1.5	10.5
11	Pulmonary hemorrhage originating in the perinatal period (P26)	28	1.4	9.8
12	Sudden infant death syndrome (R95)	24	1.2	8.4
13	Newborn affected by other complications of labor and delivery (P03)	17	0.8	*
14	Newborn affected by maternal hypertensive disorders (P00.0)	13	0.6	*
14	Interstitial emphysema and related conditions originating in the perinatal period (P25)	13	0.6	*
...	All other causes (Residual)	331	16.2	115.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2011

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Hispanic, both sexes				
...	All causes	1,451	100.0	158.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	362	24.9	39.4
2	Sudden infant death syndrome (R95)	237	16.3	25.8
3	Accidents (unintentional injuries) (V01-X59)	143	9.9	15.6
4	Diseases of the circulatory system (I00-I99)	85	5.9	9.3
5	Assault (homicide) (*U01,X85-Y09)	49	3.4	5.3
6	Diarrhea and gastroenteritis of infectious origin (A09)	38	2.6	4.1
7	Septicemia (A40-A41)	33	2.3	3.6
8	Influenza and pneumonia (J09-J18)	30	2.1	3.3
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	18	1.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	1.1	*
11	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	14	1.0	*
11	Necrotizing enterocolitis of newborn (P77)	14	1.0	*
13	Chronic respiratory disease originating in the perinatal period (P27)	13	0.9	*
14	Malignant neoplasms (C00-C97)	12	0.8	*
15	Anoxic brain damage, not elsewhere classified (G93.1)	8	0.6	*
15	Renal failure and other disorders of kidney (N17-N19,N25,N27)	8	0.6	*
...	All other causes (Residual)	371	25.6	40.4
Postneonates, Hispanic, male				
...	All causes	805	100.0	172.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	176	21.9	37.6
2	Sudden infant death syndrome (R95)	137	17.0	29.3
3	Accidents (unintentional injuries) (V01-X59)	80	9.9	17.1
4	Diseases of the circulatory system (I00-I99)	48	6.0	10.3
5	Assault (homicide) (*U01,X85-Y09)	30	3.7	6.4
6	Diarrhea and gastroenteritis of infectious origin (A09)	23	2.9	4.9
7	Influenza and pneumonia (J09-J18)	21	2.6	4.5
8	Septicemia (A40-A41)	15	1.9	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.4	*
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	10	1.2	*
10	Chronic respiratory disease originating in the perinatal period (P27)	10	1.2	*
12	Necrotizing enterocolitis of newborn (P77)	7	0.9	*
13	Malignant neoplasms (C00-C97)	6	0.7	*
13	Anoxic brain damage, not elsewhere classified (G93.1)	6	0.7	*
15	Meningitis (G00,G03)	5	0.6	*
...	All other causes (Residual)	220	27.3	47.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2011

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Hispanic, female				
...	All causes	646	100.0	143.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	186	28.8	41.3
2	Sudden infant death syndrome (R95)	100	15.5	22.2
3	Accidents (unintentional injuries) (V01-X59)	63	9.8	14.0
4	Diseases of the circulatory system (I00-I99)	37	5.7	8.2
5	Assault (homicide) (*U01,X85-Y09)	19	2.9	*
6	Septicemia (A40-A41)	18	2.8	*
7	Diarrhea and gastroenteritis of infectious origin (A09)	15	2.3	*
8	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	10	1.5	*
9	Influenza and pneumonia (J09-J18)	9	1.4	*
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	8	1.2	*
11	Necrotizing enterocolitis of newborn (P77)	7	1.1	*
12	Malignant neoplasms (C00-C97)	6	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.8	*
14	Renal failure and other disorders of kidney (N17-N19,N25,N27)	4	0.6	*
15	Acute bronchitis and acute bronchiolitis (J20-J21)	3	0.5	*
15	Chronic respiratory disease originating in the perinatal period (P27)	3	0.5	*
15	Bacterial sepsis of newborn (P36)	3	0.5	*
...	All other causes (Residual)	150	23.2	33.3
Postneonates, Non-Hispanic, white, both sexes				
...	All causes	3,666	100.0	170.8
1	Sudden infant death syndrome (R95)	899	24.5	41.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	701	19.1	32.7
3	Accidents (unintentional injuries) (V01-X59)	501	13.7	23.3
4	Diseases of the circulatory system (I00-I99)	150	4.1	7.0
5	Assault (homicide) (*U01,X85-Y09)	104	2.8	4.8
6	Influenza and pneumonia (J09-J18)	91	2.5	4.2
7	Diarrhea and gastroenteritis of infectious origin (A09)	87	2.4	4.1
8	Septicemia (A40-A41)	62	1.7	2.9
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	39	1.1	1.8
10	Chronic respiratory disease originating in the perinatal period (P27)	34	0.9	1.6
11	Malignant neoplasms (C00-C97)	27	0.7	1.3
11	Anoxic brain damage, not elsewhere classified (G93.1)	27	0.7	1.3
13	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	25	0.7	1.2
14	Acute bronchitis and acute bronchiolitis (J20-J21)	19	0.5	*
15	Meningitis (G00,G03)	15	0.4	*
15	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	15	0.4	*
...	All other causes (Residual)	870	23.7	40.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2011

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Non-Hispanic, white, male				
...	All causes	2,097	100.0	190.3
1	Sudden infant death syndrome (R95)	529	25.2	48.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	355	16.9	32.2
3	Accidents (unintentional injuries) (V01-X59)	285	13.6	25.9
4	Diseases of the circulatory system (I00-I99)	84	4.0	7.6
5	Assault (homicide) (*U01,X85-Y09)	59	2.8	5.4
6	Influenza and pneumonia (J09-J18)	55	2.6	5.0
7	Diarrhea and gastroenteritis of infectious origin (A09)	48	2.3	4.4
8	Septicemia (A40-A41)	29	1.4	2.6
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	21	1.0	1.9
10	Anoxic brain damage, not elsewhere classified (G93.1)	20	1.0	1.8
11	Chronic respiratory disease originating in the perinatal period (P27)	18	0.9	*
12	Malignant neoplasms (C00-C97)	17	0.8	*
13	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	13	0.6	*
14	Acute bronchitis and acute bronchiolitis (J20-J21)	11	0.5	*
14	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	11	0.5	*
...	All other causes (Residual)	542	25.8	49.2
Postneonates, Non-Hispanic, white, female				
...	All causes	1,569	100.0	150.2
1	Sudden infant death syndrome (R95)	370	23.6	35.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	346	22.1	33.1
3	Accidents (unintentional injuries) (V01-X59)	216	13.8	20.7
4	Diseases of the circulatory system (I00-I99)	66	4.2	6.3
5	Assault (homicide) (*U01,X85-Y09)	45	2.9	4.3
6	Diarrhea and gastroenteritis of infectious origin (A09)	39	2.5	3.7
7	Influenza and pneumonia (J09-J18)	36	2.3	3.4
8	Septicemia (A40-A41)	33	2.1	3.2
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	18	1.1	*
10	Chronic respiratory disease originating in the perinatal period (P27)	16	1.0	*
11	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	12	0.8	*
12	Malignant neoplasms (C00-C97)	10	0.6	*
13	Acute bronchitis and acute bronchiolitis (J20-J21)	8	0.5	*
13	Necrotizing enterocolitis of newborn (P77)	8	0.5	*
15	Anoxic brain damage, not elsewhere classified (G93.1)	7	0.4	*
...	All other causes (Residual)	339	21.6	32.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2011

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Non-Hispanic, black, both sexes				
...	All causes	2,410	100.0	413.8
1	Sudden infant death syndrome (R95)	524	21.7	90.0
2	Accidents (unintentional injuries) (V01-X59)	337	14.0	57.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	320	13.3	55.0
4	Diseases of the circulatory system (I00-I99)	126	5.2	21.6
5	Diarrhea and gastroenteritis of infectious origin (A09)	116	4.8	19.9
6	Assault (homicide) (*U01,X85-Y09)	88	3.7	15.1
7	Influenza and pneumonia (J09-J18)	80	3.3	13.7
8	Septicemia (A40-A41)	57	2.4	9.8
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	47	2.0	8.1
10	Chronic respiratory disease originating in the perinatal period (P27)	39	1.6	6.7
11	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	17	0.7	*
11	Necrotizing enterocolitis of newborn (P77)	17	0.7	*
13	Meningitis (G00,G03)	16	0.7	*
14	Acute bronchitis and acute bronchiolitis (J20-J21)	14	0.6	*
15	Malignant neoplasms (C00-C97)	11	0.5	*
15	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	11	0.5	*
15	Renal failure and other disorders of kidney (N17-N19,N25,N27)	11	0.5	*
...	All other causes (Residual)	579	24.0	99.4
Postneonates, Non-Hispanic, black, male				
...	All causes	1,360	100.0	459.5
1	Sudden infant death syndrome (R95)	303	22.3	102.4
2	Accidents (unintentional injuries) (V01-X59)	183	13.5	61.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	165	12.1	55.8
4	Diarrhea and gastroenteritis of infectious origin (A09)	71	5.2	24.0
5	Diseases of the circulatory system (I00-I99)	66	4.9	22.3
6	Assault (homicide) (*U01,X85-Y09)	52	3.8	17.6
7	Influenza and pneumonia (J09-J18)	45	3.3	15.2
8	Septicemia (A40-A41)	35	2.6	11.8
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	31	2.3	10.5
10	Chronic respiratory disease originating in the perinatal period (P27)	19	1.4	*
11	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	13	1.0	*
12	Acute bronchitis and acute bronchiolitis (J20-J21)	9	0.7	*
13	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	8	0.6	*
14	Malignant neoplasms (C00-C97)	7	0.5	*
14	Meningitis (G00,G03)	7	0.5	*
14	Necrotizing enterocolitis of newborn (P77)	7	0.5	*
...	All other causes (Residual)	339	24.9	114.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2011

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Non-Hispanic, black, female				
...	All causes	1,050	100.0	366.6
1	Sudden infant death syndrome (R95)	221	21.0	77.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	155	14.8	54.1
3	Accidents (unintentional injuries) (V01-X59)	154	14.7	53.8
4	Diseases of the circulatory system (I00-I99)	60	5.7	20.9
5	Diarrhea and gastroenteritis of infectious origin (A09)	45	4.3	15.7
6	Assault (homicide) (*U01,X85-Y09)	36	3.4	12.6
7	Influenza and pneumonia (J09-J18)	35	3.3	12.2
8	Septicemia (A40-A41)	22	2.1	7.7
9	Chronic respiratory disease originating in the perinatal period (P27)	20	1.9	7.0
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	16	1.5	*
11	Necrotizing enterocolitis of newborn (P77)	10	1.0	*
12	Meningitis (G00,G03)	9	0.9	*
13	Renal failure and other disorders of kidney (N17-N19,N25,N27)	6	0.6	*
14	Acute bronchitis and acute bronchiolitis (J20-J21)	5	0.5	*
14	Newborn affected by complications of placenta, cord and membranes (P02)	5	0.5	*
...	All other causes (Residual)	251	23.9	87.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011