

RESEARCH BRIEF

Understanding Barriers to Cervical Cancer Screening in Women With Access to Care, Behavioral Risk Factor Surveillance System, 2014

Anatasha Crawford, PhD, MPH; Vicki Benard, PhD; Jessica King, MPH;
Cheryl C. Thomas, MSPH

Suggested citation for this article: Crawford A, Benard V, King J, Thomas CC. Understanding Barriers to Cervical Cancer Screening in Women With Access to Care, Behavioral Risk Factor Surveillance System, 2014. *Prev Chronic Dis* 2016;13:160225. DOI: <https://doi.org/10.5888/pcd13.160225>.

PEER REVIEWED

Abstract

Cervical cancer screening can save lives when abnormal cervical lesions and early cancers are detected and treated; however, many women are not screened as recommended. We used the Behavioral Risk Factor Surveillance System survey to examine nonfinancial barriers to cervical cancer screening among women who reported having insurance and a personal doctor or health care provider. Among these women, a higher proportion who were never or rarely screened reported having multiple chronic conditions. The results of this study underscore the importance of incorporating preventive clinical services into the management of one or more chronic conditions.

Objective

Widespread use of the Papanicolaou (Pap) test has decreased cervical cancer incidence and deaths. Over half of all new cervical cancers are estimated to occur in women who have never or rarely been screened (1). Limited or no access to health care is a known barrier to screening (2). In a 2012 study of women with no cervical cancer screening in the past 5 years, nearly 70% had health insurance and a regular doctor or health care provider (3). This analysis examines nonfinancial barriers to meeting cervical

screening recommendations, focusing on women aged 40 to 65 years who may be seeking other preventive screening (eg, mammogram, colonoscopy) (4).

Methods

The Behavioral Risk Factor Surveillance System (BRFSS) survey is a state-based, random-digit-dialed telephone survey of the civilian, noninstitutionalized adult population of the United States (5). Survey data were available for all 50 states and the District of Columbia in 2014.

Women were asked if they had ever had a Pap test and if so, when this test was last performed. We selected women who were aged 40 to 65 years, reported having medical insurance and at least one personal doctor or health care provider, had not had a hysterectomy, and were not pregnant at the time of the survey. In addition to screening with a Pap test alone every 3 years, current cervical cancer screening recommendations include the use of the human papillomavirus (HPV) test with the Pap test every 5 years (4). Because HPV testing rates could not be assessed for all 50 states and the District of Columbia, respondents were categorized as never or rarely screened if they reported never having a Pap test or not having one in more than 5 years to account for the possibility that a woman may have had an HPV test (women aged 30 to 65 years who want to lengthen the screening interval and be screened with a combination of Pap test and HPV test every 5 years). Comparisons were made with women who were up to date with screening (Pap test within 3 years). Respondents who refused to answer or answered “don't know or not sure” were excluded. Eleven chronic conditions (heart attack, heart disease, stroke, asthma, skin cancer, cancer other than skin, chronic obstructive pulmonary disease [COPD], arthritis, depression, kidney disease, and diabetes) were ascertained by BRFSS and analyzed for this study. BRFSS data were weighted by using advanced raking techniques (6) and were



The opinions expressed by authors contributing to this journal do not necessarily reflect the opinions of the U.S. Department of Health and Human Services, the Public Health Service, the Centers for Disease Control and Prevention, or the authors' affiliated institutions.

age-adjusted to the 2014 BRFSS population. We performed χ^2 testing to compare the characteristics of the respondents across screening history (screened versus never or rarely screened).

Results

All variables were significant (except “asthma now” and cancer other than skin”) when comparing screened versus never or rarely screened women ($P < .05$) (Table). Compared with women screened on time, a higher proportion of women never or rarely screened were aged 60 to 65 years (27.6%), Asian/Pacific Islander (7.0%), never married (16.2%), obese (37.3%), current smokers (25.6%), and had an annual income less than \$10,000 (11%). A significantly higher proportion of preventive care measures (mammogram [83.2%], clinical breast examination [73.6%], and colorectal cancer screening [69.9%]) were observed among women who received timely cervical cancer screening when compared with women never or rarely screened ($P = .001$).

Women who were never or rarely screened for cervical cancer had a higher prevalence of ever reporting 1 of 7 chronic conditions: heart disease (4.9%), COPD (13.7%), arthritis (38.1%), depression (31.4%), kidney disease (3.8%), or diabetes (15.4%) than women who were regularly screened ($P < .01$). Higher proportions of never or rarely screened women also reported having had a heart attack (4.2%) or a stroke (4.5%) in their lifetime than regularly screened women. Women with skin cancer were more likely to be screened for cervical cancer (6.1%, $P = .004$); however, there was no significant difference ($P = .05$) in the proportions of women screened for cervical cancer who had had other forms of cancer. A significantly higher proportion of women ($P = .001$) who were never or rarely screened had more than 1 or 2 chronic conditions (3 or 4 chronic conditions, 16.6%; 5 or more chronic conditions, 3.8%) (Figure). Women screened every 3 years were more likely to have no chronic conditions (48.0%, $P = .001$).




Figure. Percentage of women aged 40 to 65 years with health insurance and a regular health care provider screened for cervical cancer by Papanicolaou (Pap) test, by number of chronic diseases, BRFSS, 2014. Chronic diseases analyzed were heart attack, heart disease, stroke, asthma, chronic obstructive pulmonary disease, arthritis, depression, kidney disease, diabetes, skin cancer, and cancer other than skin. Women who had had a hysterectomy or were pregnant at the time of the survey were excluded. Data were age-adjusted to the 2014 BRFSS population. Screening (once every 3 years) is based on the US Preventive Services Task Force recommendations for cervical cancer screening. Since HPV testing could not be assessed for all 50 states and the District of Columbia, “on time” is based on having had a Pap test within the past 3 years. Never or rarely screened refers to women who ever had a Pap test in more than 5 years to account for the possibility that a woman may have had an HPV test (women aged 30 to 65 years who want to lengthen the screening interval can be screened with a combination of Pap test and HPV test every 5 years). Data for women screened more than 3 years ago but less than 5 years ago are not shown. Abbreviation: BRFSS, Behavioral Risk Factor Surveillance System; HPV, human papilloma virus.

Discussion

When we examined factors related to cervical cancer screening among women who reported having health insurance and access to a regular doctor or health care provider, a larger proportion of women with multiple chronic conditions reported not receiving the recommended screening for cervical cancer. Our findings were similar to others indicating that insured women with arthritis, diabetes, and myocardial infarction were less likely to be screened for cervical cancer (7–10). In addition, we found that a larger proportion of women with COPD, depression, heart disease, or kidney disease did not adhere to cervical cancer screening recommendations compared with women without these conditions. Studies of women with chronic conditions who received regular health care from primary care physicians and specialists showed similar findings when examining breast and colorectal cancer screening (8,11).

Limitations of this study are not examining younger women who were eligible for screening and using self-reported survey data. Previous studies suggest that age, lack of awareness, lack of transportation, fatalistic health beliefs, low health literacy, poor patient compliance to provider recommendations, and low-quality health

care services may prevent insured women with chronic conditions from obtaining timely cervical cancer screening (3,7,12). The primary reason women adhere to timely cancer screening is because of encouragement from a provider; however, disease management for women with multiple chronic conditions is given greater priority than disease prevention (7,9). Additional research is recommended to determine if physicians can effectively balance managing patients' chronic conditions and ensuring that patients receive recommended preventive care services.

Acknowledgments

This research was supported in part by the appointment of M.L.S. to the Research Participation Program at the Centers for Disease Control and Prevention (CDC) administered by the Oak Ridge Institute for Science and Education through an interagency agreement between the US Department of Energy and CDC. The findings and conclusions in this article are those of the authors and do not necessarily represent the official position of CDC.

Author Information

Corresponding Author: Anatasha Crawford, PhD, MPH, Centers for Disease Control and Prevention, Division of Cancer Prevention and Control, Epidemiology and Applied Research Branch, 4770 Buford Hwy, Atlanta, GA 30341. Telephone: 770-488-1089. Email: wwp6@cdc.gov.

Author Affiliations: Vicki Benard, Jessica King, Cheryll C. Thomas, Centers for Disease Control and Prevention, Division of Cancer Prevention and Control, Atlanta, Georgia.

References

1. Leyden WA, Manos MM, Geiger AM, Weinmann S, Mouchawar J, Bischoff K, et al. Cervical cancer in women with comprehensive health care access: attributable factors in the screening process. *J Natl Cancer Inst* 2005;97(9):675–83.
2. Freeman HWB. Excess cervical cancer mortality: a marker for low access to health care in poor communities. Rockville (MD): National Cancer Institute, Center to Reduce Cancer Health Disparities; 2005.
3. Benard VB, Thomas CC, King J, Massetti GM, Doria-Rose VP, Saraiya M. Vital signs: cervical cancer incidence, mortality, and screening — United States, 2007–2012. *MMWR Morb Mortal Wkly Rep* 2014;63(44):1004–9.
4. Centers for Disease Control and Prevention. Cervical cancer screening guidelines for average-risk women. <http://www.cdc.gov/cancer/cervical/pdf/guidelines.pdf>. Accessed April 10, 2016.
5. Centers for Disease Control and Prevention. Behavioral Risk Factor Surveillance System. 2015. <http://www.cdc.gov/brfss/>. Accessed April 10, 2016.
6. Centers for Disease Control and Prevention. Methodologic changes in the Behavioral Risk Factor Surveillance System in 2011 and potential effects on prevalence estimates. *MMWR Morb Mortal Wkly Rep* 2012;61(22):410–3.
7. Guo F, Hirth JM, Berenson AB. Effects of cardiovascular disease on compliance with cervical and breast cancer screening recommendations among adult women. *J Womens Health (Larchmt)* 2015;24(8):641–7.
8. Liu BY, O'Malley J, Mori M, Fagnan LJ, Lieberman D, Morris CD, et al. The association of type and number of chronic diseases with breast, cervical, and colorectal cancer screening. *J Am Board Fam Med* 2014;27(5):669–81.
9. Marshall JG, Cowell JM, Campbell ES, McNaughton DB. Regional variations in cancer screening rates found in women with diabetes. *Nurs Res* 2010;59(1):34–41.
10. Owens MD, Beckles GL, Ho KK, Gorrell P, Brady J, Kaftarian JS. Women with diagnosed diabetes across the life stages: underuse of recommended preventive care services. *J Womens Health (Larchmt)* 2008;17(9):1415–23.
11. Kiefe CI, Funkhouser E, Fouad MN, May DS. Chronic disease as a barrier to breast and cervical cancer screening. *J Gen Intern Med* 1998;13(6):357–65.
12. Akers AY, Newmann SJ, Smith JS. Factors underlying disparities in cervical cancer incidence, screening, and treatment in the United States. *Curr Probl Cancer* 2007; 31(3):157–81.

Table

Table. Demographic Characteristics of US Women^a Aged 40–65 Years With Health Insurance and a Regular Health Care Provider (N = 66,402) Tested for Cervical Cancer by Papanicolaou (Pap) Test, BRFSS, 2014

| Characteristic | Screened on Time ^b | Never or Rarely Screened ^b | P Value ^d |
|-----------------------------------|-------------------------------|---------------------------------------|----------------------|
| | N (%) ^c | | |
| Demographic Characteristic | | | |
| Age, y | | | |
| Total | 60,371 (100.0) | 6,031 (100.0) | .001 |
| 40–44 | 9,095 (21.9) | 577 (16.0) | |
| 45–49 | 9,731 (17.9) | 709 (14.1) | |
| 50–54 | 12,138 (22.9) | 1,122 (21.6) | |
| 55–59 | 12,993 (17.8) | 1,391 (20.7) | |
| 60–65 | 16,414 (19.4) | 2,232 (27.6) | |
| Race/ethnicity | | | |
| Total | 59,618 (100.0) | 5,938 (100.0) | .02 |
| Non-Hispanic white | 48,771 (71.4) | 4,803 (71.4) | |
| Non-Hispanic black | 4,963 (11.7) | 413 (10.1) | |
| Asian/Pacific Islander | 1,069 (4.6) | 175 (7.0) | |
| American Indian/Alaskan Native | 727 (0.8) | 124 (1.2) | |
| Other | 1,134 (1.4) | 146 (1.6) | |
| Hispanic | 2,954 (10.0) | 277 (8.7) | |
| Marital status | | | |
| Total | 60,017 (100.0) | 5,983 (100.0) | .001 |
| Married | 38,834 (68.0) | 2,878 (49.5) | |
| Divorced | 9,581 (13.6) | 1,174 (18.9) | |
| Widowed | 3,473 (4.4) | 655 (9.2) | |
| Separated | 1,356 (2.7) | 179 (3.8) | |
| Never married | 5,622 (9.1) | 983 (16.2) | |
| Member of an unmarried couple | 1,151 (2.3) | 114 (2.4) | |
| Annual income, \$ | | | |

Abbreviation: BRFSS, Behavioral Risk Factor Surveillance System.

^a Women who had not had a hysterectomy and are not currently pregnant.

^b Adapted from the US Preventive Services Task Force recommendations for cervical cancer. Since human papilloma virus testing could not be assessed for all 50 states and the District of Columbia, “on time” is based on having had a Pap test within the past 3 years. “Never or rarely screened” is based on ever having had a Pap test in more than 5 years to account for the possibility that a woman may have had a human papilloma virus test. Data for women screened more than 3 years but less than 5 years ago is not shown.

^c Percentages are age-adjusted to the 2014 BRFSS population.

^d χ^2 Testing used to determine association.

^e Metropolitan is defined as the center city of a metropolitan statistical area or outside the center city of a metropolitan statistical area but does not include the county.

^f Current smoker is defined as smoked at least 100 cigarettes in their lifetime and now smoke every day or some days.

^g Mammogram within the past 2 years.

^h Fecal occult blood test (FOBT) in 1 year, colonoscopy in 10 years or flexible sigmoidoscopy in 5 years and FOBT in 3 years.

(continued on next page)

The opinions expressed by authors contributing to this journal do not necessarily reflect the opinions of the U.S. Department of Health and Human Services, the Public Health Service, the Centers for Disease Control and Prevention, or the authors’ affiliated institutions.

(continued)

Table. Demographic Characteristics of US Women^a Aged 40–65 Years With Health Insurance and a Regular Health Care Provider (N = 66,402) Tested for Cervical Cancer by Papanicolaou (Pap) Test, BRFSS, 2014

| Characteristic | Screened on Time ^b | Never or Rarely Screened ^b | P Value ^d |
|--|-------------------------------|---------------------------------------|----------------------|
| | N (%) ^c | | |
| Total | 53,727 (100.0) | 5,247 (100.0) | .001 |
| <10,000 | 1,837 (4.0) | 461 (11.0) | |
| 10,000–14,999 | 1,806 (3.6) | 464 (10.2) | |
| 15,000–19,999 | 2,269 (4.6) | 471 (9.1) | |
| 20,000–24,999 | 2,894 (5.4) | 545 (10.0) | |
| 25,000–34,999 | 4,068 (7.8) | 633 (12.4) | |
| 35,000–49,999 | 7,062 (12.6) | 808 (12.9) | |
| 50,000–74,999 | 9,887 (16.8) | 758 (13.0) | |
| ≥75,000 | 23,904 (45.3) | 1,107 (21.4) | |
| Education | | | |
| Total | 60,230 (100.0) | 6,006 (100.0) | .001 |
| Did not graduate high school | 2,109 (7.7) | 521 (15.0) | |
| Graduated high school | 12,477 (22.3) | 1,920 (33.0) | |
| Attended college/technical school | 16,250 (32.3) | 1,753 (31.6) | |
| Graduated college/technical school | 29,394 (37.7) | 1,812 (20.4) | |
| Body mass index, kg/m² | | | |
| Total | 56,054 (100.0) | 5,457 (100.0) | .001 |
| Underweight, ≤18 | 845 (1.6) | 133 (2.2) | |
| Normal, ≤25 | 21,645 (38.0) | 1,670 (32.1) | |
| Overweight, >25 to <30 | 17,264 (30.5) | 1,531 (28.3) | |
| Obese, ≥30 | 16,300 (29.9) | 2,123 (37.3) | |
| Residence | | | |
| Total | 42,245 (100.0) | 4,274 (100.0) | .003 |
| Metropolitan ^e | 22,885 (65.5) | 1,938 (59.2) | |
| Nonmetropolitan | 19,360 (34.5) | 2,336 (40.8) | |
| Smoking status | | | |
| Total | 60,127 (100.0) | 6,010 (100.0) | .001 |

Abbreviation: BRFSS, Behavioral Risk Factor Surveillance System.

^a Women who had not had a hysterectomy and are not currently pregnant.

^b Adapted from the US Preventive Services Task Force recommendations for cervical cancer. Since human papilloma virus testing could not be assessed for all 50 states and the District of Columbia, “on time” is based on having had a Pap test within the past 3 years. “Never or rarely screened” is based on ever having had a Pap test in more than 5 years to account for the possibility that a woman may have had a human papilloma virus test. Data for women screened more than 3 years but less than 5 years ago is not shown.

^c Percentages are age-adjusted to the 2014 BRFSS population.

^d χ^2 Testing used to determine association.

^e Metropolitan is defined as the center city of a metropolitan statistical area or outside the center city of a metropolitan statistical area but does not include the county.

^f Current smoker is defined as smoked at least 100 cigarettes in their lifetime and now smoke every day or some days.

^g Mammogram within the past 2 years.

^h Fecal occult blood test (FOBT) in 1 year, colonoscopy in 10 years or flexible sigmoidoscopy in 5 years and FOBT in 3 years.

(continued on next page)

The opinions expressed by authors contributing to this journal do not necessarily reflect the opinions of the U.S. Department of Health and Human Services, the Public Health Service, the Centers for Disease Control and Prevention, or the authors’ affiliated institutions.

(continued)

Table. Demographic Characteristics of US Women^a Aged 40–65 Years With Health Insurance and a Regular Health Care Provider (N = 66,402) Tested for Cervical Cancer by Papanicolaou (Pap) Test, BRFSS, 2014

| Characteristic | Screened on Time ^b | Never or Rarely Screened ^b | P Value ^d |
|--|-------------------------------|---------------------------------------|----------------------|
| | N (%) ^c | | |
| Current smoker ^f | 7,477 (12.8) | 1,492 (25.6) | |
| Former smoker | 14,790 (23.3) | 1,386 (23.6) | |
| Never smoked | 37,860 (63.9) | 3,132 (50.8) | |
| Use of Preventive Care | | | |
| Breast cancer screening | | | |
| Total | 60,200 (100.0) | 5,980 (100.0) | .001 |
| Yes ^g | 50,717 (83.2) | 1,895 (34.5) | |
| No | 9,483 (16.8) | 4,085 (65.5) | |
| Clinical breast examination | | | |
| Total | 60,037 (100.0) | 5,913 (100.0) | .001 |
| In last year | 44,531 (73.6) | 1,159 (22.0) | |
| >1 Year ago | 14,159 (23.4) | 3,627 (56.5) | |
| Never | 1,347 (3.0) | 1,127 (21.5) | |
| Colorectal cancer screening, aged 50–65 y^h | | | |
| Total | 40,214 (100.0) | 4,558 (100.0) | .001 |
| Yes | 28,993 (69.9) | 1,643 (34.5) | |
| No | 11,221 (30.1) | 2,915 (65.5) | |
| Chronic Disease History | | | |
| Heart attack | | | |
| Total | 60,258 (100.0) | 5,990 (100.0) | .001 |
| Yes | 1,216 (2.0) | 316 (4.2) | |
| No | 59,042 (98) | 5,674 (95.8) | |
| Heart disease | | | |
| Total | 60,173 (100.0) | 5,979 (100.0) | .001 |
| Yes | 1,436 (2.4) | 331 (4.9) | |
| No | 58,737 (97.6) | 5,648 (95.1) | |
| Stroke | | | |

Abbreviation: BRFSS, Behavioral Risk Factor Surveillance System.

^a Women who had not had a hysterectomy and are not currently pregnant.

^b Adapted from the US Preventive Services Task Force recommendations for cervical cancer. Since human papilloma virus testing could not be assessed for all 50 states and the District of Columbia, “on time” is based on having had a Pap test within the past 3 years. “Never or rarely screened” is based on ever having had a Pap test in more than 5 years to account for the possibility that a woman may have had a human papilloma virus test. Data for women screened more than 3 years but less than 5 years ago is not shown.

^c Percentages are age-adjusted to the 2014 BRFSS population.

^d χ^2 Testing used to determine association.

^e Metropolitan is defined as the center city of a metropolitan statistical area or outside the center city of a metropolitan statistical area but does not include the county.

^f Current smoker is defined as smoked at least 100 cigarettes in their lifetime and now smoke every day or some days.

^g Mammogram within the past 2 years.

^h Fecal occult blood test (FOBT) in 1 year, colonoscopy in 10 years or flexible sigmoidoscopy in 5 years and FOBT in 3 years.

(continued on next page)

The opinions expressed by authors contributing to this journal do not necessarily reflect the opinions of the U.S. Department of Health and Human Services, the Public Health Service, the Centers for Disease Control and Prevention, or the authors’ affiliated institutions.

(continued)

Table. Demographic Characteristics of US Women^a Aged 40–65 Years With Health Insurance and a Regular Health Care Provider (N = 66,402) Tested for Cervical Cancer by Papanicolaou (Pap) Test, BRFSS, 2014

| Characteristic | Screened on Time ^b | Never or Rarely Screened ^b | P Value ^d |
|--|-------------------------------|---------------------------------------|----------------------|
| | N (%) ^c | | |
| Total | 60,259 (100.0) | 6,011 (100.0) | .001 |
| Yes | 1,327 (2.1) | 290 (4.5) | |
| No | 58,932 (97.9) | 5,721 (95.5) | |
| Asthma | | | |
| Total | 60,225 (100.0) | 6,016 (100.0) | .004 |
| Yes | 8,626 (14.3) | 1,085 (18.1) | |
| No | 51,599 (85.7) | 4,931 (81.9) | |
| Asthma now | | | |
| Total | 8,441 (100.0) | 1,063 (100.0) | .26 |
| Yes | 6,468 (76.8) | 851 (79.6) | |
| No | 1,973 (23.2) | 212 (20.4) | |
| Chronic obstructive pulmonary disease | | | |
| Total | 60,191 (100.0) | 6,004 (100.0) | .001 |
| Yes | 3,645 (6.0) | 834 (13.7) | |
| No | 56,546 (94.0) | 5,170 (86.3) | |
| Arthritis | | | |
| Total | 60,113 (100.0) | 6,000 (100.0) | .001 |
| Yes | 19,462 (31.4) | 2,405 (38.1) | |
| No | 40,651 (68.6) | 3,595 (61.9) | |
| Depression | | | |
| Total | 60,147 (100.0) | 6,001 (100.0) | .001 |
| Yes | 14,142 (22.0) | 1,810 (31.4) | |
| No | 46,005 (78.0) | 4,191 (68.6) | |
| Kidney disease | | | |
| Total | 60,251 (100.0) | 6,002 (100.0) | .001 |
| Yes | 1,501 (2.5) | 241 (3.8) | |
| No | 58,750 (97.5) | 5,761 (96.2) | |

Abbreviation: BRFSS, Behavioral Risk Factor Surveillance System.

^a Women who had not had a hysterectomy and are not currently pregnant.

^b Adapted from the US Preventive Services Task Force recommendations for cervical cancer. Since human papilloma virus testing could not be assessed for all 50 states and the District of Columbia, “on time” is based on having had a Pap test within the past 3 years. “Never or rarely screened” is based on ever having had a Pap test in more than 5 years to account for the possibility that a woman may have had a human papilloma virus test. Data for women screened more than 3 years but less than 5 years ago is not shown.

^c Percentages are age-adjusted to the 2014 BRFSS population.

^d χ^2 Testing used to determine association.

^e Metropolitan is defined as the center city of a metropolitan statistical area or outside the center city of a metropolitan statistical area but does not include the county.

^f Current smoker is defined as smoked at least 100 cigarettes in their lifetime and now smoke every day or some days.

^g Mammogram within the past 2 years.

^h Fecal occult blood test (FOBT) in 1 year, colonoscopy in 10 years or flexible sigmoidoscopy in 5 years and FOBT in 3 years.

(continued on next page)

The opinions expressed by authors contributing to this journal do not necessarily reflect the opinions of the U.S. Department of Health and Human Services, the Public Health Service, the Centers for Disease Control and Prevention, or the authors' affiliated institutions.

(continued)

Table. Demographic Characteristics of US Women^a Aged 40–65 Years With Health Insurance and a Regular Health Care Provider (N = 66,402) Tested for Cervical Cancer by Papanicolaou (Pap) Test, BRFSS, 2014

| Characteristic | Screened on Time ^b | Never or Rarely Screened ^b | P Value ^d |
|------------------------|-------------------------------|---------------------------------------|----------------------|
| | N (%) ^c | | |
| Diabetes | | | |
| Total | 60,302 (100.0) | 6,019 (100.0) | .001 |
| Yes | 5,652 (9.6) | 1,011 (15.4) | |
| Only during pregnancy | 1,061 (1.9) | 99 (1.5) | |
| No | 52,714 (86.8) | 4,804 (81.4) | |
| Prediabetes | 875 (1.7) | 105 (1.7) | |
| Skin cancer | | | |
| Total | 60,261 (100.0) | 6,014 (100.0) | .004 |
| Yes | 4,095 (6.1) | 288 (4.4) | |
| No | 56,166 (93.9) | 5,726 (95.6) | |
| Cancer, nonskin | | | |
| Total | 60,251 (100.0) | 6,003 (100.0) | .05 |
| Yes | 4,288 (6.6) | 402 (5.4) | |
| No | 55,963 (93.4) | 5,601 (94.6) | |

Abbreviation: BRFSS, Behavioral Risk Factor Surveillance System.

^a Women who had not had a hysterectomy and are not currently pregnant.

^b Adapted from the US Preventive Services Task Force recommendations for cervical cancer. Since human papilloma virus testing could not be assessed for all 50 states and the District of Columbia, “on time” is based on having had a Pap test within the past 3 years. “Never or rarely screened” is based on ever having had a Pap test in more than 5 years to account for the possibility that a woman may have had a human papilloma virus test. Data for women screened more than 3 years but less than 5 years ago is not shown.

^c Percentages are age-adjusted to the 2014 BRFSS population.

^d χ^2 Testing used to determine association.

^e Metropolitan is defined as the center city of a metropolitan statistical area or outside the center city of a metropolitan statistical area but does not include the county.

^f Current smoker is defined as smoked at least 100 cigarettes in their lifetime and now smoke every day or some days.

^g Mammogram within the past 2 years.

^h Fecal occult blood test (FOBT) in 1 year, colonoscopy in 10 years or flexible sigmoidoscopy in 5 years and FOBT in 3 years.

The opinions expressed by authors contributing to this journal do not necessarily reflect the opinions of the U.S. Department of Health and Human Services, the Public Health Service, the Centers for Disease Control and Prevention, or the authors' affiliated institutions.